

Chromatographic and immunological identification of GnRH (Gonadotropin-releasing hormone) variants. Occurrence of mammalian and a salmon-like GnRH in the forebrain of an eutherian mammal : Hydrochaeris hydrochaeris (Mammalia, Rodentia

Item Type	Article
Authors	Montaner, A.D.;Somoza, G.M.;King, A.J.;Bianchinni, J.J.;Blis, C.G.;Affani, J.M.
Citation	MONTANER AD, SOMOZA GM, KING JA, BIANCHINI JJ, BOLIS CG, AFFANNI JM. Chromatographic and immunological identification of GnRH (Gonadotropin-releasing hormone) variants. Occurrence of mammalian and a salmon-like GnRH in the forebrain of an eutherian mammal#: Hydrochaeris hydrochaeris (Mammalia, Rodentia). Regulatory peptides [Internet]. 1998
Publisher	Elsevier
Journal	Regulatory peptides
Rights	Attribution 3.0 United States
Download date	2024-04-24 19:16:18
Item License	http://creativecommons.org/licenses/by/3.0/us/
Link to Item	https://infospace.mrc.ac.za/handle/11288/595216

Chromatographic and immunological identification of GnRH (Gonadotropin-releasing hormone) variants. Occurrence of mammalian and a salmon-like GnRH in the forebrain of an eutherian mammal: *Hydrochaeris hydrochaeris* (Mammalia, Rodentia).

Alejandro D. Montaner^{a,b}, Gustavo M. Somoza^{a,*}, Judy A. King^c, Juan J. Bianchini^a,
C. Giuliana Bolis^d, Jorge M. Affanni^{a,b}

^aInstituto de Neurociencia (INEUCI-CONICET), Departamento de Biología, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires (UBA), Pabellón II, Ciudad Universitaria, 1428 Buenos Aires, Argentina

^bCentro de Investigaciones Científicas y Transferencia de Tecnología a la Producción (CONICET), Matteri y España 3105, Diamante, Entre Ríos, Argentina

^cMedical Research Council Regulatory Peptides Research Unit, Department of Chemical Pathology, University of Cape Town Medical School and Groote Schuur Hospital, Observatory 7925, Cape Town, South Africa

^dBiología Generale, Istituto di Scienze Farmacologiche, Facoltà di Farmacia, Università di Milano, Via Balzaretti 9-20133 Milan, Italy

Received 25 June 1997; received in revised form 22 December 1997; accepted 29 December 1997

Abstract

The molecular variants of Gonadotropin releasing hormone (GnRH) in brain extracts of the eutherian mammal *Hydrochaeris hydrochaeris* (Mammalia, Rodentia) were characterized. An indirect method combining reverse-phase high-performance liquid chromatography (RP-HPLC) and radioimmunoassay (RIA) with different antisera was used. Two different forebrain regions (olfactory bulbs and preoptic–hypothalamic region) were analyzed. Characterization of RP-HPLC fractions from preoptic–hypothalamic extracts with three different RIA systems revealed two immunoreactive GnRH (ir-GnRH) peaks coeluting with mammalian GnRH (mGnRH) and salmon GnRH (sGnRH) synthetic standards. These results were additionally supported by serial dilution studies with specific antisera. Similar results were obtained from olfactory bulb extracts with the same methodology. However, a third ir-GnRH peak in a similar position to that of chicken GnRH II (cIIGnRH) synthetic standard was revealed. As far as we know, this is the first report showing chromatographic and immunological evidences for the presence of a second GnRH variant in the forebrain of an eutherian mammal. © 1998 Elsevier Science B.V.

Keywords: Gonadotropin-releasing hormone; GnRH; HPLC; Radioimmunoassay; Mammalian; Capybara

1. Introduction

GnRH, originally isolated from ovine and porcine brains, is the hypothalamic hormone that regulates the reproductive system by stimulation of gonadotropins from the anterior pituitary gland [1]. This hormone was considered to be a unique molecular form, until GnRH molecular

variants were demonstrated. To date, nine molecular variants have been characterized in vertebrates [2,3] and two other forms were recently characterized from a protochordate [4]. They all are decapeptides, with a very conserved structure: amino-terminus (pGlu) and carboxy-terminus (NH₂) are modified and conserved, all of them having common positions in 1, 2, 4, 9 and 10.

cIIGnRH is expressed together with mGnRH in the brain of most of metatherian species studied [5–7]. Most of the eutherian species studied as yet have only mGnRH,

*Corresponding author: Tel./fax: +54 1 7818016; e-mail: somoza@biolo.bg.fcen.uba.ar

Table 1
Radioimmunoassay characteristics

Antiserum	¹²⁵ I-GnRH	STD	Titer	Percentage of cross-reactivity				
				mGnRH	cIGnRH	cIIGnRH	sGnRH	IIGnRH
m1076	mGnRH	mGnRH	1:50 000	100	59	0.4	0.3	< 0.01
EL-15	mGnRH	mGnRH	1:175 000	100	< 0.01	< 0.01	< 0.01	0.05
cII678	mGnRH	mGnRH	1:40 000	100	416	81	473	3.1
cII675	cIIGnRH	cIIGnRH	1:40 000	< 0.01	< 0.01	100	< 0.01	< 0.01
sI668	sGnRH	sGnRH	1:10 000	< 0.01	< 0.01	0.7	100	0.4
sGnRH#2	sGnRH	sGnRH	1:300 000	< 0.01	< 0.01	1.58	100	0.08
PBL#45	sGnRH	sGnRH	1:250 000	100	133	25.9	100	24
PBL#49	sGnRH	sGnRH	1:150 000	93.3	333	32.6	100	2.0

PBL#45, PBL#49 and CII678 antisera were used in the screening assays. The following antisera were used in homologous assays: m1076, EL-15, CII675, S1668 and anti sGnRH#2. ¹²⁵I-GnRH: radioiodinated hormone; STD: synthetic standard; IIGnRH: lamprey GnRH I. The percentages of crossreactivities of the different RIA systems used were reported by Dr. King, Dr. Okuzawa [24] and Dr. Yu [19], and retested in our conditions. The percentages of crossreactivities of the RIA system with EL-15 were calculated in our laboratory.

with the order Insectivora excepted [3]. mGnRH is coexpressed with cIIGnRH [7,9,10] in the brain of different species of this primitive order [8]. However, there are few reports concerning identification of GnRH variants in eutherian mammal species other than Insectivora [11–13].

The aim of this study was to determine the nature of GnRH variants in the brain of the amphibian mammal *Hydrochaeris hydrochaeris*. It is commonly named ‘capybara’, ‘carpincho’ or water hog. This is the largest living rodent and can not be confused with any other form [14]. It may be 1.25 m (50 in.) long and weight 50 kg or more. It is shy and associates in groups along the banks of lakes and rivers of South America. It swims and dives readily and commonly enters water to elude predators. It is strictly vegetarian. The female bears a single litter of three to eight young each year and gestation takes about 100 to 120 days [15].

This model bears special interest because of the following reasons: this animal belongs to the suborder Hystricomorpha considered to be one of the oldest groups of rodents [16]. It is the only living species belonging to the family Hydrochaeridae. This group underwent great radiation in America during Pliocene and Pleistocene periods [16]. Its anatomical features are similar to those of the oldest extinguished rodents [16,17]. Some authors consider this group the mother group of all remaining rodents [18]. Its origin is found in fauna from African Eocene, members from which, migrated across the South Atlantic to South America by rafts [19]. However there are no data about brain physiology except the paper of Affanni et al. [20].

2. Materials and methods

2.1. Animals

Two adult males and one adult female, weighing among 20–50 kg, of *Hydrochaeris hydrochaeris* were used. They were captured by licensed animal officers in the surrounding areas of Diamante (Province of Entre Ríos, Argentina).

They were immediately taken to the laboratory. They were anesthetized by a combination of ketamine hydrochloride (40 mg/kg, i.m.) and sodium thiopental (60 mg/kg, i.m.). The brains were removed, and brain regions were obtained by dissection. The pieces were immediately frozen in dry ice, pooled and stored at –70°C until extraction of peptides.

2.2. Tissue extraction

Preoptic–hypothalamic region (4.5 g) and olfactory bulbs (1.5 g) were independently homogenized in acetone: 1 M HCl (100:3, v/v) at 4°C with an Ultraturrax homogenizer. The extraction mixtures were stirred at 4°C and filtered through Whatman N°1 filter paper. The insoluble material was reextracted in acetone: 0.01 M HCl in 40% of the original volume, stirred for 5 min and refiltered. The combined filtrates were treated with petroleum ether (b.p. 30–60°C) for five successive times as previously described [21,22]. The final aqueous phases were then concentrated to less than 1 ml using a vacuum concentrator.

2.3. Reverse phase-high performance liquid chromatography

The concentrated extracts were filtered through a 0.45 µm Millipore HA filter. They were injected through 1 ml injection loop onto a Lichrospher 100 RP18 column (4 × 250 mm; 5 µm particle size, Merck) with a C18 guard column.

A Konic liquid chromatograph was used. It was programmed according Sherwood et al. [21]. The flow rate was kept at ml/min. The filtrate was injected at the beginning of a 10 min period of 17% acetonitrile in TEAF (0.25 M formic acid adjusted with triethylamine to pH 6.5). The acetonitrile concentration was then increased linearly over a 7 min period to 24%. The elution continued under isocratic conditions for a further 53 min.

Blank chromatographs injecting 1 ml of TEAF were performed before each RP-HPLC of tissue extracts to

check possible contamination of the system. The fractions were radioimmunoassayed under the same conditions as tissue extracts. Fresh standards of GnRH (200 ng of each one) were pooled and diluted in TEAF. They were injected together and chromatographed after the tissue extracts chromatography. Mammalian GnRH, chicken GnRH I (cIGnRH), cIIGnRH and sGnRH were obtained from Peninsula Laboratories Inc. (San Carlos, CA, USA).

RP-HPLC collected fractions were lyophilized and resuspended in RIA buffer. Ir-GnRH was tested using different GnRH RIA systems.

2.4. Radioimmunoassays

Radioimmunoassays were made using synthetic mGnRH, cIIGnRH and sGnRH as some of us previously described [23–25]. The different GnRH standards were radiolabeled following established protocols [26] with minor modifications [23]. Extracts of preoptic–hypothalamic region and olfactory bulbs were analyzed for ir-GnRH in a first screening. PBL#45, PBL#49 and cII678 assays were used. See Table 1 for RIA systems and antisera characteristics.

Fractions with ir-GnRH suspected to be caused by an endogenous determined variant were pooled and then analyzed with specific RIAs. The slopes of the displacement curves were compared with those of synthetic standards by ANOVA.

3. Results

The elution positions of the different GnRH synthetic standards were: mGnRH, between 20–22 min; cIGnRH, 22–24 min; cIIGnRH, 27–29 min. The more hydrophobic sGnRH eluted between 50–60 min depending upon the chromatography.

Three different RIA systems were used as a first screening (see Table 1).

3.1. Preoptic-hypothalamic region

RP-HPLC fractions of extracts from this region revealed two main ir-GnRH peaks when analyzed with the three RIA systems (Fig. 1a, b and c). The first ir-GnRH eluting peak (fractions 22–24) coeluted with mGnRH synthetic standard. The second one (fractions 60–63) coeluted with synthetic sGnRH.

Fractions showing ir-mGnRH and ir-sGnRH were pooled independently and serial dilution displacements were performed. Fractions 22–24, which coeluted with synthetic mGnRH, were then tested with two homologous mGnRH RIA systems using m1076 and E-15 antisera (Fig. 2a and b). The slopes of serial dilutions of pooled fractions were not significantly different from those of synthetic standards. The same analysis was performed in fractions




Fig. 1. RP-HPLC of extracts from preoptic–hypothalamic region analyzed with RIAs with the following antisera: (a) PBL #45, (b) PBL #49 and (c) cII 678. Arrows indicate the elution position of four different GnRH standards. The left arrow shows the elution position of mGnRH. The second arrow represents cIGnRH, the third is cIIGnRH, and the last one is sGnRH.

coeluting with synthetic sGnRH. These fractions were analyzed with two specific sGnRH RIA systems (s1668 and sGnRH#2 antisera). Both yielded curves parallel to those of sGnRH synthetic standard (Fig. 3a and b).




Fig. 2. Parallelism of serial dilutions of fractions 22–24 from RP-HPLC extract from preoptic–hypothalamic region in comparison with synthetic mGnRH. (a) mGnRH 1076 and (b) EL-15 antisera. The slopes of both curves were not significantly different from those of synthetic standard.

Retention time and quantification of immunoreactive GnRH peaks are summarized in Table 2.

3.2. Olfactory bulbs

A similar screening was performed to characterize ir-GnRH fractions in these structures. Three ir-GnRH peaks were revealed (Fig. 4a, b and c). The first one (fractions 21–23) eluted in the same position as mGnRH synthetic. The second ir-GnRH peak (fractions 27–28), coeluted with cIIgNnRH synthetic standard and the last one eluted in a position similar to that of sGnRH (fractions 49–50).

Unfortunately ir-GnRH levels in all ir-GnRH fractions were low and serial displacement studies were not so clear. Nevertheless the three ir-GnRH fractions appeared displacing radioiodinated GnRH in homologous assays (EL-15, cII675 and s1668 RIA systems, data not shown).

Retention times and quantification of ir-GnRH peaks of olfactory bulbs extracts are showed in Table 3.




Fig. 3. Parallelism of serial dilutions of fractions 60–62 from HPLC extract of preoptic–hypothalamic region with respect to synthetic sGnRH, using (a) sGnRH 1668 antiserum and (b) sGnRH#2 antiserum. The slopes of both curves were not significantly different from those of synthetic standard.

4. Discussion

The analysis of preoptic–hypothalamic region extract from ‘capybara’ revealed two ir-GnRH peaks. These peaks have chromatographic and immunological features resembling those of mGnRH and sGnRH synthetic standards.

Characterization of RP-HPLC fractions using: cII678, PBL #45 and PBL #49, revealed two ir-GnRH peaks coeluting with mGnRH and sGnRH synthetic standards. Parallelism of serial fractional dilution showing ir-mGnRH and ir-sGnRH to standards curves was seen in each case using specific antisera (mGnRH 1076 and EL-15 antisera for mGnRH and sGnRH 1668 and sGnRH#2 antisera for sGnRH). In both cases (mGnRH and sGnRH assays) serial dilution of pooled fractions with specific antisera reinforced RP-HPLC and RIA data.

Similar results were obtained with extract from olfactory bulbs. Three ir-GnRH peaks were observed by RIA with polyspecific antisera. However serial dilution experiments

Table 2
Retention time and quantification of ir-GnRH peaks in preoptic–hypothalamic extract after RP-HPLC with different antisera.

Tissue	Peak	Ret. time (min)	ir-GnRH (pg/100 μ l) ^a				Ret. time GnRH-std ^b
			M1076	EL-15	S1668	sGnRH#2	
PHR	I	22–24	70	130	–	–	22 (M)
	II	60–63	–	–	20	10	60 (S)

^a Immunoreactive GnRH in pooled fractions from each peak from extract of preoptic–hypothalamic region (PHR).

^b Retention times of synthetic peptides which were run immediately after the biological tissue runs. (M), mGnRH; (S), sGnRH.




Fig. 4. RP-HPLC of the olfactory bulbs extract with (a) PBL #45, (b) PBL #49 and (c) cII 678 antisera. Arrows show the position of four different GnRH variants. See Fig. 1 for references.

with these ir-GnRH material were not convincing. This uncertainty was probably due to low GnRH levels. On the other hand, the presence of a third ir-GnRH peak in an elution position similar to that of cII GnRH should be confirmed by further experiments.

Our results show that similarly to other mammals the dominant form of GnRH in adult ‘capybara’ is mGnRH. In mammals and other vertebrates, neurons expressing the GnRH molecular variant related to gonadotropin secretion arise from the olfactory placode [28–32]. These neurons are distributed through the terminal nerve, the medial–septal, the preoptic area and the rostral hypothalamus. These neurons belong to the terminal nerve-septo-preoptic system [28,33]. This system is related to GnRH release to the pituitary portal system acting on gonadotropic cells of the adenohypophysis [1,28]. On the other hand, cII GnRH was shown together with mGnRH in several species of metatherian and early evolved eutherian mammals [5–7,9,10]. In these animals, olfactory regions and areas of the ventral forebrain contain neurons and fibres immunoreactive to mGnRH. There is also a large population of ir-cII GnRH cells in the midbrain [3,9,10].

Muske [28] proposed that gnathostomic vertebrates have two main GnRH systems with different embryonic origins. They are: a) the anterior, terminal nerve-septo-preoptic system, which is the main regulator of gonadotropin release and b) the posterior system formed by neurons in periventricular regions of the posterior diencephalon and/or midbrain.

During vertebrate evolution, the conservation of mesencephalic cII GnRH is well known [3]. This form has been reported in cartilaginous fish [31], bony fish [22,23,25,27,34–36], amphibians [37–39], reptiles [40,41], birds [42–44], most of metatherian mammals [5–7] and early evolved eutherian species [7,9,10].

The above mentioned data suggest that there was a gene duplication in the earliest vertebrates. It is important to note that the conservation of structure and neuroanatomical location of cII GnRH neurons would imply a very important nongonadotropic role of this variant. However the biological meaning and function of this form remain to be discovered [28]. Our data with extracts from olfactory bulbs would suggest that cII GnRH is also being expressed in this mammal. More assays are needed to confirm these data. Moreover, further studies on extracts from mesencephalic region must be performed.

Table 3
Retention time and quantification of ir-GnRH peaks in the extract of olfactory bulbs chromatographed on RP-HPLC.

Tissue		Ret. Time (min)	ir-GnRH (pg/100 μ l) ^a					Ret. time GnRH-std ^b
Olfactory	Peak		M1076	EL-15	cII675	S1668	sGnRH#2	
Bulbs	I	21–23	10	10	–	–	–	22 (M)
	II	27–28	–	–	5	–	–	27 (CII)
	III	50–51	–	–	–	4	2	50 (S)

^a Immunoreactive GnRH in pooled fractions from each peak (corresponding to the retention time shown in Fig. 4).

^b Retention times of synthetic GnRH peptides. (M), mammalian GnRH; (CII), chicken GnRH II; (S), salmon GnRH.

Our data show the surprising fact that there are two coexisting GnRH variants in the anterior GnRH system of 'capybara'. These data suggest the following possibilities: a) both variants would be involved in the regulation of adenohipophyseal hormones; b) one of them, probably mGnRH would related to gonadotropin secretion whereas the other variant is expressed by nerve terminals ending in the pars nervosa; and c) one variant acts on pituitary regulation and the other would serve as a neurotransmitter or neuromodulator. Regarding these possibilities, we must point out that several forms of GnRH also stimulate the secretion of growth hormone in teleost fish [45–47]. Moreover, some variants have a differential gonadotropin-growth hormone-releasing activity [48]. This effect was recently shown in rats [49]. These findings support the idea of GnRH acting on two adenohipophyseal hormones.

There is also a worth mentioning fact. This is represented by the demonstration of ir-GnRH fibres entering the neural lobe in amphibians [50,51], mammals [52,53] and even in humans [52]. Even more, it appears that extension of hypophysiotropic neurosecretory systems (GnRH, GHRH and somatostatin) into the pituitary pars nervosa is characteristic of many mammalian species [53]. From that structure they could be released directly to the bloodstream and/or be involved to paracrine interactions within the neurohypophysis.

In our case, the presence of ir-sGnRH in extracts from preoptic–hypothalamic region opens a series of evolutionary questions. These questions are not only concerned with physiological aspects and differential actions of GnRH variants but with the fact that sGnRH has only been found in teleost fish [3]. Such an unusual presence provides abundant material for speculation.

New experiments must be done to characterize GnRH variants in mesencephalic extracts from 'capybara'. The neuroanatomical location of ir-mGnRH and s-GnRH cell bodies and fibres must also be determined.

Although final conclusions need isolation, purification and sequencing of the molecules, these results offer encouraging possibilities of further advances.

As far as we know, this is the first report showing immunological and chromatographic evidences for the presence of two GnRH molecular variants: mGnRH and a

salmon GnRH-like form, in the forebrain of an eutherian mammal.

Acknowledgements

This work was supported by Grants from CONICET (National Research Council of Argentina) for promoting the Centro de Investigaciones Científicas y Transferencia de Tecnología a la Producción (Diamante, Entre Ríos) to Dr. J.M.A. and the Medical Research Council of South Africa and the University of Cape Town to Dr. J.A.K. We are indebted to Dr. Hernán Aldana-Marcos and Dr. Omar Iódice for their help in the management of animals. We would also like to thank Prof. José Osinaldi from the Dirección General de Desarrollo Agrícola y Recursos Naturales (Gobierno de la Provincia de Entre Ríos, Secretaría de Estado de la Producción)

References

- [1] Fink G. Gonadotropin secretion and its control. In: Knobil E, Neill J, editors. The physiology of reproduction. New York: Raven Press, 1988:1349–77.
- [2] Powell JFF, Zohar Y, Elizur A et al. Three forms of gonadotropin-releasing hormone characterized from brains of one species. Proc Natl Acad Sci USA 1994;91:12081–5.
- [3] King JA, Millar RP. Evolutionary aspects of gonadotropin-releasing hormone and its receptor. Cell Mol Neurobiol 1995;15:5–23.
- [4] Powell JFF, Reska-Skinner SM, Om Prakash M et al. Two new forms of gonadotropin-releasing hormone in a protochordate and the evolutionary implications. Proc Natl Acad Sci USA 1996;93:10461–4.
- [5] King JA, Mehl AEI, Tyndale-Biscoe CH et al. A second form of GnRH, with chicken GnRH II-like properties, occurs together with mammalian GnRH in marsupial brains. Endocrinology 1989;125:2244–52.
- [6] King JA, Hinds LA, Mehl AEI et al. Chicken GnRH II occurs together with mammalian GnRH in South American species of marsupial (*Monodelphis domestica*). Peptides 1990;11:521–5.
- [7] King JA, Steneveld AA, Curlew JD et al. Identification of chicken GnRH II in the brains of metatherian and early-evolved eutherian species of mammals. Regul Pept 1994;54:467–77.
- [8] Eisenber JF. The history of mammals. In: Turner JF, editor. The mammalian radiations. An analysis of trends in evolution, adapta-

- tion, and behavior. Chicago: The University of Chicago Press, 1981:7–12.
- [9] Dellovade T, King JA, Millar RP et al. Presence and differential distribution of distinct forms of immunoreactive gonadotropin-releasing hormone in the musk shrew brain. *Neuroendocrinology* 1993;58:166–77.
- [10] Kasten TL, White SA, Norton TT et al. Characterization of two new preproGnRH mRNAs in the Tree Shrew: first direct evidence for mesencephalic GnRH gene expression in a placental mammal. *Gen Comp Endocrinol* 1996;104:7–19.
- [11] King JA, Hassan MF, Mehl AEI et al. Gonadotropin-releasing hormone molecular forms in mammalian hypothalamus. *Endocrinology* 1988;122:2742–52.
- [12] Kelsall R, Coe IR, Sherwood NM. Phylogeny and ontogeny of gonadotropin-releasing hormone: comparison of guinea pig, rat and a protochordate. *Gen Comp Endocrinol* 1990;78:479–94.
- [13] Gautron JP, Pattou E, Bauer K et al. (Hydroxyproline⁹) luteinizing hormone-releasing hormone: A novel peptide in mammalian and frog hypothalamus. *Neurochem Int* 1991;18:221–35.
- [14] Redford KH, Eisenberg JF. Mammals of the Neotropics. The Southern Cone. Volume 2. Chile, Argentina, Uruguay, Paraguay. Chicago, London: The University of Chicago Press, 1992:343–4.
- [15] Ojasti J. Estudio biológico del chigüire o capibara, Fondo Nacional de Investigaciones Agropecuarias. Caracas, Venezuela: Editorial Sucre, 1973:275.
- [16] Patterson B, Pascual R. The fossil mammals fauna of South America. In: Keast A, Erk FC, Glass B, editors. *Evolution, Mammals and Southern Continents*. New York: The University of New York Press, Albany, 1972:247–309.
- [17] Hershkovitz E. The recent mammals of the neotropical region: a zoogeographic and ecological review. *Quart Rev Biol* 1969;44:1–70.
- [18] Novasek MJ. Mammalian phylogeny: Shaking the tree. *Nature* 1992;365:121–5.
- [19] Lavocat L. What is an hystricomorph. *Symp Zool Soc Lond* 1974;34:7–20.
- [20] Affanni JM, Scaravilli AM, Gori AM et al. La actividad eléctrica del neocortex de la vizcacha (*Lagostomus maximus maximus*), de la chinchilla (*Chinchilla brevicaudata*) y del carpincho (*Hydrochoerus hydrochaeris*) (Mammalia Rodentia). *Physis* 1973;C32:181–90.
- [21] Sherwood N, Zoeller RT, Moore FL. Multiple forms of gonadotropin-releasing hormone in amphibians brain. *Gen Comp Endocrinol* 1986;61:313–22.
- [22] Yu KL, Sherwood NM, Peter RE. Differential distribution of two molecular forms of gonadotropin-releasing hormone in discrete brain areas of goldfish (*Carassius auratus*). *Peptides* 1988;9:625–30.
- [23] Somoza GM, Stefano AV, D'Eramo JL et al. Immunoreactive GnRH suggesting a third variant form of GnRH in addition to cIIIGnRH and sGnRH in the brain and pituitary gland of *Prochilodus lineatus* (Characiformes). *Gen Comp Endocrinol* 1994;94:44–52.
- [24] Somoza GM, Paz DA, Stefano AV et al. Identification of immunoreactive mammalian gonadotropin-releasing hormone in the brain of metamorphic larvae of *Bufo arenarum* Hensel (Amphibia: Anura). *Int J Dev Neurosci* 1996;14:663–72.
- [25] Stefano AV, Fridman O, Somoza GM. Chromatographic and immunological evidence for a third form of GnRH in addition to cIIIGnRH and sGnRH in the brain of *Odontesthes bonariensis*. In: Goetz F, Thomas P, editors. *Reproductive Physiology of Fish*, The University of Texas at Austin, Marine Science Institute, TX, USA, 1995:79.
- [26] Yu KL, Nahoraniak CS, Peter RE et al. Brain distribution of radioimmunoassayable gonadotropin-releasing hormone in female goldfish: seasonal variation and periovulatory changes. *Gen Comp Endocrinol* 1987;67:234–46.
- [27] Okuzawa K, Amano M, Kobayashi M et al. Differences in sGnRH and cGnRH-II contents in discrete brain areas of male and female rainbow trout according to age and stage of maturity. *Gen Comp Endocrinol* 1990;80:116–26.
- [28] Muske LE. Evolution of Gonadotropin-releasing hormone (GnRH) neuronal systems. *Brain Behav Evol* 1993;42:215–43.
- [29] Witkin JW, Silverman AJ. Luteinizing Hormone-Releasing Hormone (LHRH) in rat olfactory system. *J Comp Neurol* 1983;218:426–32.
- [30] Schwanzel-Fukuda M, Abraham S, Crossin KL et al. Immunocytochemical demonstration of Neural Cell Adhesion Molecule (NCAM) along the migration route of Luteinizing Hormone-Releasing Hormone (LHRH) neurons in mice. *J Comp Neurol* 1992;321:1–18.
- [31] Murakami S, Kikuyama S, Arai Y. The origin of Luteinizing Hormone-Releasing Hormone (LHRH) neurons in newts (*Cynops pyrrhogaster*): the effect of olfactory placode ablation. *Cell Tissue Res* 1992;269:21–7.
- [32] Murakami S, Seki T, Wakabayashi K et al. The ontogeny of luteinizing hormone releasing hormone (LHRH) producing neurons in the chick embryo: possible evidence for migrating LHRH neurons from the olfactory epithelium expressing a highly polysialylated neural cell adhesion molecule. *Neurosci Res* 1991;12:421–31.
- [33] Jennes L, Conn M. Gonadotropin-Releasing Hormone and its receptors in rat brain. *Front Neuroendocrinol* 1994;15:51–77.
- [34] Lovejoy DA, King JA, Sherwood NM et al. Identification of gonadotropin-releasing hormone and associated binding substances in the blood serum of a holocephalan (*Hydrolagus colliei*). *Peptides* 1993;14:1237–43.
- [35] Yamamoto N, Oka Y, Amano M et al. Multiple gonadotropin-releasing hormone (GnRH)-Immunoreactive Systems in the brain of the dwarf gourami, *Colisa lalia*: Immunohistochemistry and radioimmunoassay. *J Comp Neurol* 1995;355:354–68.
- [36] King JA, Millar RP, Vallarino M et al. Localization and characterization of gonadotropin-releasing hormones in the brain, gonads, and plasma of a dipnoi (lungfish *Protopterus annectens*). *Regul Pept* 1995;57:163–74.
- [37] King JA, Millar RP. Identification of His⁵, Trp⁷, Tyr⁸-GnRH (chicken GnRH-II) in amphibian brain. *Peptides* 1986;7:827–34.
- [38] Northcutt RG, Muske L. Multiple embryonic origins of gonadotropin-releasing hormone (GnRH) immunoreactive neurons. *Dev Brain Res* 1994;78:279–90.
- [39] Licht P, Tsai PS, Sotowska-Brochocka J. The nature and distribution of gonadotropin-releasing hormones in brains and plasma of ranid frogs. *Gen Comp Endocrinol* 1987;66:248–55.
- [40] Powell RC, Ciarcia G, Lance V et al. Identification of diverse molecular forms of GnRH in reptile brain. *Peptides* 1986;7:1101–8.
- [41] Lovejoy DA, Fischer WH, Parker DB et al. Primary structure of two forms of gonadotropin-releasing hormone (GnRH) from brains of the American alligator (*Alligator mississippiensis*). *Regul Pept* 1991;33:105–16.
- [42] Powell RC, Jach H, Millar RP et al. Identification of Gln⁸-GnRH and His⁵, Trp⁷, Tyr⁸-GnRH in the hypothalamus and extrahypothalamic brain of the ostrich (*Struthio camelus*). *Peptides* 1987;8:185–90.
- [43] Millam JR, Faris PL, Youngren OM et al. Immunohistochemical localization of chicken gonadotropin-releasing hormones I and II (cGnRH I and II) in turkey hen brain. *J Comp Neurol* 1993;333:68–82.
- [44] van Gils J, Absil P, Grauwels L et al. Distribution of luteinizing hormone-releasing hormones I and II (LHRH-I and -II) in the quail and chicken brains as demonstrated with antibodies directed against synthetic peptides. *J Comp Neurol* 1993;334:304–23.
- [45] Marchant TA, Chang JP, Nahorniak CS et al. Evidence that gonadotropin-releasing hormone also functions as a growth hormone-releasing factor in the goldfish. *Endocrinology* 1989;124:2509–18.
- [46] Lin XW, Lin HR, Peter RE. Growth hormone and gonadotropin secretion in the common carp (*Cyprinus carpio* L.): in vitro interactions of gonadotropin-releasing hormone, somatostatin, and the dopamine agonist apomorphine. *Gen Comp Endocrinol* 1993;89:62–71.

- [47] Lin HR, Lu M, Lin XW et al. Effects of gonadotropin-releasing hormone (GnRH) analogs and sex steroids on growth hormone (GH) secretion and growth in common carp (*Cyprinus carpio*) and grass carp (*Ctenopharyngodon idellus*). *Aquaculture* 1995;135:173–84.
- [48] Ngamvongchon S, Lovejoy DA, Fischer WH et al. Characterization of the primary structure of gonadotropin releasing hormone in the Thai catfish (*Clarias macrocephalus*). *Mol Cell Neurosci* 1992;3:17–22.
- [49] Penzias AS, Goodman SR, Rossi G et al. Preliminary evidence that GnRH has the properties of a growth hormone-releasing factor in vivo in the rat. *J Soc Gynecol Invest* 1995;2:623–9.
- [50] Andersen AC, Danger JM, Fasolo A et al. Immunohistochemical localization of gonadotropin-releasing-hormone-associated peptide in the brain of the frog. *J Comp Neurol* 1988;273:241–51.
- [51] Rastogi RK, Dimeglio M, Iela L. Immunoreactive luteinizing hormone-releasing hormone in the frog (*Rana esculenta*) brain: distribution pattern in the adult, seasonal changes, castration effects, and developmental aspects. *Gen Comp Endocrinol* 1990;78:444–58.
- [52] Anthony EL, King JC, Stopa EG. Immunocytochemical localization of LHRH in the median eminence, infundibular stalk, and neurohypophysis. Evidence for multiple sites of releasing hormone secretion in humans and other mammals. *Cell Tissue Res* 1984;236:5–14.
- [53] Anthony EL, Bruhn TO. Do some traditional hypophysiotropic hormones play nontraditional roles in the neurohypophysis? Implications of immunocytochemical studies. In: North WG, Moses AM, Share L, editors. *The Neurohypophysis: a window on brain function*. *Ann NY Acad Sci* 1993;689:469–72.