

Association between arterial compliance and anthropometry of children from four ethnic groups in South Africa: the Thusa bana study

Item Type	Article
Authors	Schutte, A.E.;Huisman, H.W.;Van Rooyen, J.M.;De Ridder, J.M.;Malan, N.T.
Citation	SCHUTTE AE, HUISMAN HW, VAN ROOYEN JM, DE RIDDER JH, MALAN NT. Associations between arterial compliance and anthropometry of children from four ethnic groups in South Africa: The THUSA BANA study. Blood pressure [Internet]. 2003 Jan 1
Journal	Blood Pressure
Rights	Attribution 3.0 United States
Download date	2025-07-30 16:15:53
Item License	http://creativecommons.org/licenses/by/3.0/us/
Link to Item	https://infospace.mrc.ac.za/handle/11288/595277

Associations between Arterial Compliance and Anthropometry of Children from Four Ethnic Groups in South Africa: The THUSA BANA Study

ALETTA ELISABETH SCHUTTE¹, HUGO WILLEM HUISMAN¹, JOHANNES MARTHINUS VAN ROOYEN¹, JOHANNES HENDRIK DE RIDDER² AND NICOLAAS THEODORE MALAN¹

From the ¹School for Physiology, Nutrition and Consumer Sciences, ²School for Biokinetics, Sport Science and Recreation, Potchefstroom University for Christian Higher Education, Potchefstroom, South Africa

Schutte AE, Huisman HW, van Rooyen JM, de Ridder JH, Malan NT. Associations between arterial compliance and anthropometry of children from four ethnic groups in South Africa: the THUSA BANA study. Blood Pressure 2003; 12: 97–103.

Aim: To investigate whether associations and differences exist regarding the arterial compliance, dietary intake and anthropometric parameters of children of four different ethnic groups in South Africa. *Study design:* In this study, 1244 children from four ethnic groups, aged 10–15 years, were randomly selected from five regions of the North West Province. Blood pressure was measured with a Finapres apparatus and analysed to obtain systemic arterial compliance. Measurements were done to obtain body mass index (BMI), waist-to-hip ratio and percentage body fat. Dietary intake was determined with a 24-h dietary recall questionnaire. *Results:* The black and mixed-origin subjects indicated the lowest values ($p \le 0.05$) for BMI, percentage body fat, and dietary intake, whereas the white and Indian subjects showed significantly higher values than the other ethnic groups. The white and Indian children had significantly higher arterial compliance than the black and mixed-origin children at all ages from 10 to 15 years. Compliance showed significant correlations with all anthropometric parameters. *Conclusion:* Since black and mixed-origin children at all ages from the other arterial compliance of undernutrition and stunted growth, it is suggested that parental undernutrition and inadequate nutrition in early life, associated with lower arterial compliance, ethnicity, *hypertension, race, undernutrition.*

INTRODUCTION

Ethnic differences in the prevalence of adult hypertension are well known [1]. The African-American adults are known to have higher blood pressure than Hispanic or non-Hispanic white adults [1], whereas Mexican American adults do not have an increased prevalence of hypertension [2], although they have a higher prevalence of cardiovascular risk factors than non-Hispanic whites [3].

Conflicting results have been reported in the paediatric literature regarding ethnic differences in blood pressure levels in children and adolescents [4–6]. Although clinical hypertension occurs less frequently in children than in adults [7–9], ample evidence supports the concept that the roots of essential hypertension extend back to childhood. Of particular importance is the documentation that elevated blood pressure in childhood often correlates with hypertension in early adulthood, thereby supporting the need to track blood pressure in children [10]. Studies of blood pressure in African-American, Hispanic and non-Hispanic children showed no consistent, statistically significant differences in blood pressure levels among

© 2003 Taylor & Francis on licence from Blood Pressure. *ISSN 0803-7051* DOI 10.1080/08037050310001075

Indians, 17.2% in whites [14], 22.2% in blacks [15] and 30.2% in the mixed-origin group [16]. However, very little is known regarding blood pressure in children of the

children and adolescents, regardless of adjustment for weight or use of auscultatory or oscillometric method

[3, 5], but there are also studies that indicated opposite

in 2000 - comprising of 77.6% Africans (black), 2.5%

Asians (Indian), 8.7% mixed origin and 10.4% whites

(European) [12] - clear differences in the prevalence of

hypertension have been observed. It has been indicated

that food poverty rates were highest among households

with the worst socio-economic conditions, headed by

Africans, followed by people of mixed origin, Indians and

whites [13]. These different ethnic groups indicated

prevalences of hypertension that varied from 14.2% in

In the South African population of just over 43 million

results [4, 11].

various ethnic groups of South Africa. Although blood pressure is the most frequently measured parameter of the peripheral vasculature, other properties such as arterial compliance, may be a more subtle index of vascular dysfunction associated with diseases such as hypertension [17]. Vascular compliance is defined as the change in volume of the artery per unit of pressure $(\Delta V/\Delta P)$ [18]. Arterial compliance can also be estimated from the simpler approach of stroke volume divided by pulse pressure [19] and it reflects both the arterial capacity and viscoelastic properties of the arterial wall. Arterial compliance is influenced by various conditions. It has been shown that with aging, compliance of large arteries decreases [20]. It has also been shown that severe obesity in children is associated with arterial wall stiffness (low arterial compliance) [21]. Other anthropometric factors, such as body size (stature), are also strongly associated with arterial compliance [22].

According to the World Health Organization [23], there is very little direct evidence about the determinants of common cardiovascular diseases in populations of sub-Saharan Africa. The aim of this study was therefore to investigate whether associations and differences exist regarding the cardiovascular and anthropometric parameters and dietary intake of children of four different ethnic groups. This study forms part of the larger THUSA BANA study (Transition and Health during Urbanization in South Africa in Children; Bana: the Setswana word for Children), which was designed to assess the health status of children in the North West Province of South Africa.

METHODS

Study design

This was an epidemiological, cross-sectional study. Forty-four schools were randomly selected from a list of schools in the five regions of the North West Province of South Africa. These schools were visited during the weeks preceding the collection of data, in order to obtain permission from the relevant school principals as well as from the parents of the children. Children within the schools were also randomly selected from class lists. Data collection took place during normal school hours.

Subjects

A total of 1244 apparently healthy children between 10 and 15 years of age were recruited from the 44 schools over a period of 2 years (2000–2001). The subjects consisted of black, white, Indian and mixed-origin children of both sexes. Equal percentages of younger and older subjects were included in each of the four race groups. Because of the magnitude of the project, all subjects were not necessarily subjected to all measurements.

The Ethics Committee of the Potchefstroom University for Christian Higher Education has approved the study and the study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki. All parents of the THUSA BANA subjects gave informed consent.

Data collection and measurements

The subjects were all introduced to the experimental setup, after which each one was separately subjected to the following procedures:

Cardiovascular parameters

The subjects were connected to a Finapres (finger-arterial pressure) apparatus [24, 25] and blood pressure was recorded continuously. After a period of rest of at least 10 min, resting blood pressure values were obtained. Blood pressure was regarded as resting when the systolic blood pressure did not change with more than 10 mmHg during the last minute of this period, otherwise the resting period was extended. The resting blood pressure was then recorded continuously for 1 min. The data was stored on magnetic tape by means of a Kyowa RTP-50A fourchannel data recorder and digitized for further analysis by means of the Fast Modelflow software program [26]. The Modelflow method digitally computes an aortic-flow waveform from a peripheral arterial pressure signal [26]. It uses a non-linear three-element model (Windkessel) of the aortic input impedance consisting of aortic characteristic impedance, arterial compliance and peripheral vascular resistance [26]. In this way the systolic (SBP), and diastolic blood pressure (DBP), and arterial compliance (C) were obtained.

The vascular unloading technique of Penáz together with the Physiocal criteria of Wesseling provided reliable, non-invasive and continuous estimates of blood pressure [25, 27]. This technique is thus an alternative to the invasive intra-arterial measurements, without the risks and ethical questions inherent to invasive measurements. Since the pressure waveform is available continuously, computations provide further information on the dynamics of the cardiovascular system on a beat-to-beat basis, similar to intra-arterial measurements [26, 28–30].

Anthropometric measurements

Anthropometric measurements were done by qualified anthropometrists under guidance of a level III anthropometrist, according to standard methods as described by Norton & Olds [31]. Maximum height was measured to the nearest 0.1 cm by means of a stadiometer with the head in the Frankfort plane. Body mass was measured to the nearest 0.1 kg by means of a calibrated electronic scale (Precision Health scale) with the subject wearing the minimum clothing. The waist and hip girths of the subjects were measured with a flexible Lufkin steel anthropometric tape to the nearest 0.1 cm. During these measurements, the subject had to stand erect with the feet together and without volitionally contracting the gluteal muscles.

The following equations were used to determine body





mass index and waist-to-hip ratio [31]: BMI = body mass (kg)/stature² (m); WHR = waist girth (cm)/hip girth (cm).

Skin-fold measurements were taken using a Harpenden Skin-fold caliper with a jaw pressure of 10 g/mm^2 and were taken to the nearest 0.2 mm by using the standard methods as described by Norton & Olds [31]. Percentage body fat was calculated according to the equation of Boileau *et al.* [32].

Dietary intake

Dietary intake data were collected by fieldworkers trained by registered dieticians. A 24-h dietary recall was collected face-to-face and the data collection interview method and nutrient coding were the same for all recalls. Food models and photo books for portion-size estimates were used for the recalls. This type of dietary assessment is widely used in international epidemiological studies [33–35]. Macronutrients (protein, fat, and carbohydrate) and fibre were calculated, and micronutrients (such as potassium, sodium and iron) and vitamins (such as folic acid) were calculated in the appropriate units, using a computer program based on the South African food composition tables [36].

Statistical analysis

All processed data was transferred to Microsoft Excel and further statistically analysed by means of the software computer package STATISTICA [37]. Due to skewed distributions all dietary variables were logarithmically transformed. Since the blood pressure and anthropometric parameters had a normal distribution, log transformations were not necessary. One-way analysis of variance (ANOVA) was used to determine whether statistical significant differences ($p \le 0.05$) existed between the different ethnical groups as far as cardiovascular and anthropometric parameters are concerned. It was followed by multiple comparisons of the group means using the Tukey honest significant difference (HSD) test for unequal group sizes [37]. The one-way analysis of covariance (ANCOVA) was used to determine significant differences ($p \le 0.05$) between the arterial compliance of the different age groups, correcting for height (Fig. 1).

Table I. Cardiovascular and anthropometric characteristics of the subjects

	n	SBP (mmHg)	DBP (mmHg)	C (ml/mmHg)	BMI (kg/m ²)	WHR	% BF	Body mass (kg)	Height (m)
Males									
Black	356	98.4 ± 14.5	64.1 ± 10.5	$1.07\pm0.39^{\rm a}$	$16.6\pm2.7^{\rm a}$	0.84 ± 0.07	$13.7\pm5.6^{\rm a}$	$35.4\pm10.1^{\rm a}$	$144.9\pm12.0^{\rm a}$
White	102	103.1 ± 14.5	65.4 ± 8.6	1.34 ± 0.50^{ab}	$19.0 \pm 3.8^{\mathrm{abc}}$	0.82 ± 0.08	17.3 ± 7.9^{ab}	47.5 ± 13.6^{abc}	$157.0 \pm 1.4^{\rm ab}$
Mixed origin	98	98.6 ± 12.5	63.5 ± 9.2	$0.99\pm0.33^{\mathrm{b}}$	$16.3 \pm 2.7^{\mathrm{b}}$	0.83 ± 0.04	14.0 ± 5.3^{b}	$34.2\pm8.3^{\rm b}$	144.1 ± 1.2^{bc}
Indian	43	95.6 ± 14.4	61.7 ± 9.4	1.15 ± 0.33	$17.0 \pm 3.3^{\circ}$	0.82 ± 0.04	15.8 ± 5.8	$39.5\pm11.0^{\rm c}$	$151.4\pm10.9^{\rm c}$
Females									
Black	414	103.8 ± 15.4	66.6 ± 11.1	$1.00\pm0.37^{\rm a}$	18.1 ± 3.9	0.78 ± 0.07	22.1 ± 7.0	$39.4 \pm 11.7^{\rm a}$	146.3 ± 11.4^{a}
White	91	105.9 ± 13.8	64.8 ± 8.2	1.19 ± 0.41^{ab}	$19.1\pm2.9^{\rm a}$	0.77 ± 0.08	$24.2\pm5.9^{\rm a}$	46.5 ± 11.5^{ab}	154.9 ± 11.2^{ab}
Mixed origin	99	103.6 ± 14.4	65.5 ± 9.1	$0.96\pm0.37^{\mathrm{b}}$	$17.2\pm3.1^{\rm a}$	0.77 ± 0.05	21.3 ± 6.7^a	$38.0\pm10.7^{\rm b}$	147.4 ± 10.7^{b}
Indian	41	98.3 ± 14.4	63.4 ± 9.1	1.11 ± 0.35	18.0 ± 3.6	0.76 ± 0.08	24.5 ± 7.0	41.5 ± 10.2	151.5 ± 9.8

Means with the same superscript letter differ significantly ($p \le 0.05$) and all significant differences have effect sizes ≥ 0.05 . Values are mean \pm standard deviation.

n, number of subjects; SBP, systolic blood pressure; DBP, diastolic blood pressure; BMI, body mass index; WHR, waist-to-hip ratio; %BF, percentage body fat.

	Group A	Group B		
Dietary variables	(Black and mixed origin)	(Indian and white)		
Protein (g/day)	61.0 ± 0.8	64.1 ± 1.6		
Animal protein (g/day)	29.5 ± 0.6	28.3 ± 1.1		
Plant protein (g/day)	$31.3\pm0.5^{\mathrm{b}}$	35.7 ± 1.1^{b}		
Saturated fat (g/day)	17.7 ± 0.4	16.9 ± 0.7		
Fibre (g/day)	$15.2\pm0.3^{\mathrm{b}}$	$17.6\pm0.6^{\mathrm{b}}$		
Iron (mg/day)	8.4 ± 0.2	8.7 ± 0.3		
Magnesium (mg/day)	$229.4\pm3.6^{\rm a}$	$246.9\pm6.9^{\rm a}$		
Potassium (mg/day)	1625.2 ± 28.1	1683.1 ± 52.2		
Sodium (mg/day)	1585.4 ± 37.9	1644.4 ± 67.5		
Folic acid (µg/day)	$169.8\pm4.5^{\rm a}$	$186.5\pm8.8^{\rm a}$		

Table II. *Mean dietary intake of the lower (Group A) and higher socio-economic (Group B) groups*

Values are mean \pm standard error.

Significantly different: ^a $p \le 0.05$; ^b $p \le 0.001$, all significant differences have effect sizes ≥ 0.02 .

The independent *t*-test was used to determine statistically significant differences ($p \le 0.05$) between the dietary intakes of the lower and higher socio-economic groups (Groups A and B). Effect sizes (*d*) were calculated with the equation:

$$\mathbf{d} = -\mathbf{x}_1 - \mathbf{x}_2 \mathbf{S} \mathbf{D}$$

(*x* = mean; SD = maximum standard deviation) and give an indication of the practical importance of the statistical significance [38]. Since the variables were normally distributed, the Pearson correlation coefficient was used to indicate statistical significant correlations ($p \le 0.05$) between arterial compliance and the anthropometric variables.

RESULTS

The overall sample consisted of 599 male and 645 female subjects. The means and SDs of cardiovascular and anthropometric characteristics of these subjects are indicated in Table I, together with the results of the Tukey test. No significant differences were indicated for SBP and DBP between any of the groups. However, arterial compliance of the white males and females indicated significantly higher values than the black and mixed-origin groups. From this table, it is also evident that the white male subjects had significantly higher BMI and body mass than the black, mixed-origin and Indian groups. They also had a significantly higher percentage body fat and height than the black and mixed-origin children. The Indian males had a significantly higher height than the mixed-origin males. The white female subjects also indicated a significantly higher BMI and percentage body fat than the mixed-origin females. They also indicated a significantly higher body mass and height than the black and mixed-origin females. The effect sizes (standardized differences) of the significant differences are indicators of practical importance, and indicate practical results of a medium to a large effect [38]. The black and mixed-origin subjects of both sexes indicated the lowest values for body mass, height, BMI and percentage body fat.

Table II provides the mean and standard errors of dietary intake data of the subjects within two groups. Subjects with lower socio-economic status and food poverty [13] were placed in Group A (black and mixed origin). These subjects were mostly subjected to living conditions in rural areas, on farms and in informal settlements. Subjects with higher socio-economic status were placed in Group B (Indian and white) in order to determine basic dietary differences between subjects from these backgrounds. These subjects lived mostly in established townships with full access to water and electricity and in upper class suburbs. It is evident from this data that subjects from a higher socio-economic background (white and Indian) had significantly higher $(p \le 0.001)$ intakes of plant protein and fibre, as well as $(p \le 0.05)$ magnesium and folic acid than those with lower socio-economic status (black and mixed origin). The effect sizes of these significant differences were small, meaning that these results are practical to use in a direction giving capacity [38].

With the aging process from 10 to 15 years, body size increases, which leads to an increase in the arterial tree, and an increased arterial compliance [39]. It is evident

Table III. Correlations between arterial compliance and anthropometric parameters

	Body mass index	Waist-to-hip ratio	Percentage body fat	Mass	Stature
Males	r = 0.45	r = -0.20	r = 0.12	r = 0.82	r = 0.91
	n = 540	n = 540	n = 539	n = 540	n = 540
Females	p = 0.00001	p = 0.00001	p = 0.006	p = 0.00001	p = 0.00001
	r = 0.54	r = -0.41	r = 0.49	r = 0.86	r = 0.87
i emaies	n = 561	n = 560	n = 560	n = 561	n = 562
	p = 0.00001	p = 0.00001	p = 0.00001	p = 0.00001	p = 0.00001

r, Pearson correlation coefficient; n, amount of subjects; p, statistical significance.

When the arterial compliance of the total subject group was correlated with the five anthropometric variables (Table III), statistical significant correlations ($p \le 0.05$) between all anthropometric parameters and compliance were found. The compliance of the female subjects showed much stronger correlations with BMI, WHR and percentage body fat, than the males. The same phenomenon was also evident when the correlations were done within the separate ethnic groups.

DISCUSSION

Since arterial compliance consists of an arterial distensibility component but also an arterial volume component, it is reasoned that an increase in body size – as can be seen in the aging process – will be positively associated with increases in compliance [39]. However, it is not only body size influencing compliance at such a young age, since it is evident from Fig. 1 that significant differences in compliance exist between the four ethnic groups, although there has been corrected for body height.

The possibility of genetic predisposition of certain ethnic groups to present low arterial compliance cannot be ruled out, but since it is clear that compliance is strongly related to body size (Table III), the origin of lower arterial compliance might also be partially explained by nutritional deficiencies. In a review of the nutritional status of South Africans [40], one of the most important observations was that between 20% and 25% of preschoolers are stunted, and therefore suffer from chronic undernutrition. Black and mixed-origin children (lower socio-economic status) had the highest prevalences (25 and 17% respectively) of undernutrition, with rural black children being the most vulnerable group. Rural black children had low mean energy intakes and although total protein intakes seemed adequate, the quality of the protein may have been jeopardized. These results have been duplicated in this study, which showed that children of black and mixed origin had lower intakes of protein, especially plant protein, higher intakes of saturated fat and lower intakes of dietary fibre. The significantly lower intakes of fibre and folic acid in these groups are an indication that fruit and vegetable intakes are not adequate.

The parental dietary intake may also play a significant role in jeopardizing the cardiovascular health of the infant. Several lines of evidence support the thesis that it is poor or imbalanced delivery of nutrients that programmes raised blood pressure in humans. Numerous animal experiments have shown that undernutrition *in utero* leads to fetal adaptations causing persisting changes in blood pressure [41].

It is therefore proposed that a vicious circle of events might be the cause of low arterial compliance in these children. The origin of this circle is parental undernutrition, in areas of low socio-economic support, which leads to children with low birth weight. This is followed by children predisposed to hypertension, and who are also subjected to undernutrition or malnutrition, thereby promoting the development of hypertension in later life.

It is suspected that household food insecurity, poverty, and other factors dictated by socio-economic realities, are more important determinants of nutrient intakes. It is clear that low energy density of weaning foods and of the diets of rural black primary school children, coupled to a low intake of fruits, vegetables, legumes and milk by many individuals are the main deficiencies in the South African diet [40].

The strong negative association of WHR of the female subjects (Table III) indicates that a high prevalence in abdominal obesity, especially in females, is associated with low arterial compliance. The black male and female subjects not only showed the lowest arterial compliance in all age groups (Fig. 1), but also showed the highest WHR, although not statistically significantly higher than the other ethnic groups. Vorster and co-workers [40] also stated that although a quarter of South Africa's children are stunted, very high prevalences of obesity are also evident, indicating that South Africans clearly have the double burden of both under- and malnutrition. The effects of stunted growth were clearly observed in the black children, as well as the children of mixed origin, who showed a significantly lower BMI, body mass, fat percentage and height than the white and Indian children. By calculating the body surface, the differences between these ethnic groups could have possibly been observed more clearly.

Parental undernutrition and inadequate nutrition in early life, associated with lower arterial compliance, could therefore lead to the onset of adult hypertension, since it has been demonstrated by Arnett *et al.* [42] that reductions in large or small arterial compliance occur early in adulthood before the appearance of hypertension. It is therefore suggested that vascular structural changes may already begin at an early age.

We conclude therefore that although there are not yet any significant differences in systolic and diastolic blood pressure between the four ethnic groups of children, there are clear significant differences in arterial compliance. This suggests that the differences in arterial compliance could be associated with the differences in the prevalence of adult hypertension between these four ethnic groups, but further study is needed. The black and mixed-origin ethnic groups (lower socio-economic status) had the lowest arterial compliance in childhood in this study (Fig. 1) and other studies also indicated that these ethnic groups had the highest prevalence of hypertension in adulthood of 22.2% and 30.2%, respectively [15, 16]. The white and Indian groups of this study (higher socio-economic status) had the highest arterial compliance in childhood (Fig. 1), and these groups also indicated the lowest prevalence of adult hypertension of 17.2% and 14.2%, respectively [14] in South Africa. These findings suggest that low arterial compliance in childhood, associated with nutritional deficiencies and stunted growth (smaller body size), may have an association with hypertension in later life, but further study is needed to confirm this suggestion.

ACKNOWLEDGEMENTS

We would like to thank Prof. F. C. Eloff, Mrs L. Malan and Mr R. Schutte for their contributions during this study, as well as the whole THUSA BANA research team of the Potchefstroom University for CHE. We would also like to thank those who have funded this study: The SA Sugar Association, the Medical Research Council of South Africa, the Potchefstroom University for CHE, Hypertension Society of SA and Department of Trade and Industry through the THRIP system. The authors are also grateful to Prof. H. S. Steyn for his statistical consultation.

REFERENCES

- Chase HP, Garg SK, Icaza G, Carmain JA, Walravens CF, Marshall G. Twenty four hour ambulatory blood pressure monitoring in healthy young adult Anglo, Hispanic and African-American subjects. Am J Hypertens 1997; 10: 18– 23.
- McMillen M, Dridz T, Haynes S Hypertension in Mexican-Americans. HHANES, Southwest United States, 1982– 1984. Washington, DC: American Public Health Association, 1985.
- Park MK, Menard SW, Yuan C. Comparison of blood pressure in children from three ethnic groups. Am J Cardiol 2001; 87: 1305–8.
- Alpert BS, Fox ME. Racial aspects of blood pressure in children and adolescents. Pediatr Clin N Am 1993; 40: 13– 22.
- Hohn AR, Dwyer KM, Dwyer JH. Blood pressure in youth from four ethnic groups: the Pasadina Prevention Project. J Pediatr 1994; 125: 368–73.
- Webber LS, Harsha DW, Phillips GT, Srinivasan SR, Simpson JW, Berenson GS. Cardiovascular risk factors in Hispanic, white and black children: the Brooks County and Bogalusa Heart Studies. Am J Epidemiol 1991; 133: 704– 14.
- Joint National Committee. The fifth report of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure. Arch Intern Med 1993; 153: 154–83.
- Sinaiko AR, Gillum RF, Jacobs DR Jr, Sopko G, Prineas RJ. Renin–angiotensin and sympathetic nervous system activity in grade school children. Hypertension 1982; 4: 299–306.
- 9. Sinaiko AR, Gomez-Marin O, Prineas RJ. Prevalence of "significant" hypertension in junior high school-aged

children: the Children and Adolescent Blood Pressure Program. J Pediatr 1989; 114: 664–9.

- Lauer RM, Clarke WR. Childhood risk factors for adult blood pressure: the Muscatine Study. Pediatrics 1984; 84: 633–41.
- 11. Gutgesell M, Terrell G, Labarthe D. Pediatric blood pressure. Ethnic comparison in a primary care center. Hypertension 1981; 3: 39–47.
- Anon. Statistics South Africa 2000 [Online]. Available at http://www.statssa.gov.za/RELEASES/DEMOGRAP/2000/ po302.htm. Accessed 10 September 2001.
- Charlton KE, Rose D. Prevalence of household food poverty in South Africa: results from a large, nationally representative survey. Public Health Nutr 2002; 5: 383–90.
- 14. Seedat YK. Hypertension in black South Africans. J Hum Hypertens 1999; 13: 97–103.
- 15. Van Rooyen JM, Kruger HS, Huisman HW, *et al.* An epidemiological study of hypertension and its determinants in a population in transition: the THUSA study. J Hum Hypertens 2000; 14: 779–787.
- 16. Steyn K, Jooste PL, Fourie JM, Parry CD, Rossouw JE. Hypertension in the coloured population in the Cape Peninsula. SA Med J 1986; 69: 165–169.
- 17. Resnick LM, Militianu D, Cunnings AJ, *et al.* Pulse waveform analysis of arterial compliance: relation to other techniques, age, and metabolic variables. Am J Hypertens 2000; 13: 1243–9.
- Van Bortel LM, Spek JJ. Influence of aging on arterial compliance. J Hum Hypertens 1998; 12: 583–6.
- Dart AM, Kingwell BA. Pulse pressure a review of mechanisms and clinical relevance. J Am Coll Cardiol 2001; 37: 975–84.
- Benetos A, Laurent S, Hoeks AP, Boutouyrie PH, Safar ME. Arterial alterations with aging and high blood pressure. A non-invasive study of carotid and femoral arteries. Arterioscler Thromb 1993; 13: 90–7.
- 21. Tounian P, Aggoun Y, Dubern B, *et al.* Presence of increased stiffness of the common carotid artery and endothelial dysfunction in severely obese children: a prospective study. Lancet 2001; 358: 1400–4.
- 22. Soma J, Aakhus S, Angelsen BA, Skjaerpe T. Influence of body size and left ventricular ejection dynamics on total arterial compliance determined using Doppler echocardiography and subclavian artery pulse tracings in healthy humans. Blood Press 1998; 7: 239–46.
- World Health Organization. International Society of Hypertension Guidelines for the Management of Hypertension. J Hypertens 1999; 17: 151–83.
- Wesseling KH, Settels JJ, De Wit B. The measurement of continuous finger arterial pressure noninvasively in stationary subjects. In: Schmidt TH, Dembroski TM, Blümchen G, editors. Biological factors in cardiovascular disease. Berlin: Springer. 1986: 355–75.
- 25. Silke B, McAuley D. Accuracy and precision of blood pressure determination with the Finapres: an overview using re-sampling statistics. J Hum Hypertens 1998; 12: 403–9.
- 26. Wesseling KH, Jansen JRC, Settels JJ, Schreuder JJ. Computation of aortic flow from pressure in humans using a nonlinear, three-element model. J Appl Physiol 1993; 75: 2566–73.
- McAuley D, Silke B, Farrel S. Reliability of blood pressure determination with the Finapres with altered physiological states or pharmacodynamic conditions. Clin Auton Res 1997; 7: 179–84.

- 28. Langewouters GJ, Settels JJ, Roelandt R, Wesseling KH. Why use Finapres or Portapres rather than intra-arterial or intermittent non-invasive techniques of blood pressure measurement? J Med Eng Technol 1998; 22: 37–43.
- Imholz BPM, Wieling W, Van Montfrans GA, Wesseling KH. Fifteen years experience with finger arterial pressure monitoring: assessment of the technology. Cardiovasc Res 1998; 38: 605–16.
- 30. Harms MPM, Wesseling KH, Pott F, Jenstrup M, Van Goudoever J, Secher NH, *et al.* Continuous stroke volume monitoring by modeling flow from non-invasive measurement of arterial pressure in humans under orthostatic stress. Clin Sci 1999; 97: 291–301.
- Norton K, Olds T Anthropometrica: a textbook of body measurement for sports and health courses. Sydney: UNSW Press, 1996.
- Boileau RA, Lohman TG, Slaughter MH. Exercise and body composition of children and youth. Scand J Sport Sci 1985; 7: 17–27.
- Simons-Morton DG, Hunsberger SA, Van Horn L, *et al.* Nutrient intake and blood pressure in the Dietary Intervention Study in Children. Hypertension 1997; 29: 930–6.
- Falkner B, Sherif K, Michel S, Kushner H. Dietary nutrients and blood pressure in urban minority adolescents at risk for hypertension. Arch Pediatr Adolesc Med 2000; 154: 918– 22.
- Hajjar IM, Grim CE, George V, Kotchen TA. Impact of diet on blood pressure and age-related changes in blood pressure in the US population. Arch Intern Med 2001; 161: 589–93.
- Langenhoven M, Kruger M, Gouws E, Faber M MRC food composition tables. Tygerberg: Medical Research Council, 1991.
- 37. StatSoft, Inc. STATISTICA for Windows (Computer program manual). Tulsa, OK: StatSoft, Inc., 2000.

- 38. Cohen J Statistical power analysis for the behavioural sciences, 2nd edition. Hillsdale, NJ: Lawrence Erlbaum Associates, Year?.
- 39. Soma J, Aakhus S, Angelsen BA, Skjaerpe T. Influence of body size and left ventricular ejection dynamics on total arterial compliance determined using Doppler echocardiography and subclavian artery pulse tracings in healthy humans. Blood Press 1998; 7: 239–46.
- Vorster HH, Oosthuizen W, Jerling JC, Veldman FJ, Burger HM The nutritional status of South Africans. A review of the literature from 1975–1996. Durban: Health Systems Trust, 1997.
- 41. Barker DJP. In utero programming of cardiovascular disease. Theriogenology 2000; 53: 555–74.
- 42. Arnett DK, Glasser SP, McVeigh G, *et al.* Blood pressure and arterial compliance in young adults: the Minnesota children's blood pressure study. Am J Hypertens 2001; 14: 200–5.

Submitted December 7, 2002; accepted October 14, 2002

Address for correspondence:

A. E. Schutte School for Physiology, Nutrition and Consumer Science Potchefstroom University for CHE Private Bag X6001 Potchefstroom 2520 South Africa Tel: +27 18 2992435 Fax: +27 18 2992433 E-mail: flgaes@puknet.puk.ac.za Copyright © 2003 EBSCO Publishing