

Respiratory atopic disease, Ascaris-immunoglobulin E and tuberculin testing in urban South African children

Item Type	Article
Authors	Obihara, C.C.;Beyers, N.;Gie, R.P.;Hoekstra, M.O.;Fincham, J.E.;Marais, B.J.;Lombard, C.J.;Dini, L.A.;Kimpen, J.L.L.
Citation	Obihara CC, Beyers N, Gie RP, Hoekstra MO, Fincham JE, Marais BJ, et al. Respiratory atopic disease, Ascaris-immunoglobulin E and tuberculin testing in urban South African children. CLINICAL AND EXPERIMENTAL ALLERGY
Publisher	Blackwell Publishing Ltd.
Journal	Clinical and Experimental Allergy
Rights	Attribution 3.0 United States
Download date	2025-05-09 13:32:26
Item License	http://creativecommons.org/licenses/by/3.0/us/
Link to Item	https://infospace.mrc.ac.za/handle/11288/595234

ORIGINAL PAPER

Journal compilation© 2006 Blackwell Publishing Ltd

Respiratory atopic disease, *Ascaris*-immunoglobulin E and tuberculin testing in urban South African children

C. C. Obihara*[†], N. Beyers[†], R. P. Gie[†], M. O. Hoekstra^{*}, J. E. Fincham[‡], B. J. Marais[†], C. J. Lombard[§], L. A. Dini[¶] and J. L. L. Kimpen^{*}

* Wilhelmina Children's Hospital, University Medical Center, Utrecht, The Netherlands, [†]Desmond Tutu TB Centre, Department of Paediatrics and Child Health, Faculty of Health Sciences, Stellenbosch University, Stellenbosch, South Africa, [‡]Nutritional Intervention Research Unit, [§]Biostatistics Unit, Medical Research Council of South Africa, Cape Town, South Africa and [¶]Parasitology Reference Unit, National Institute for Communicable Diseases, National Health Laboratory Services, Johannesburg, South Africa

Clinical and Experimental Allergy

Summary

Background Epidemiological relation of intestinal helminth infection and atopic disease, both associated with a T-helper (Th) 2 immune response, is controversial, as it has been reported that helminth infection may either suppress or pre-dispose to atopic disease. This relation has not been tested in an area with a high burden of *Mycobacterium tuberculosis* (MTB) infection, a known Th1-stimulating infection.

Objective To study the association of intestinal helminth infection and atopic disease in a community where helminth infection is endemic and MTB infection is high.

Methods Three-hundred and fifty-nine randomly selected children aged 6–14 years from a poor urban suburb were tested with allergy questionnaire, skin prick test (SPT) to common aeroallergens, *Ascaris*-specific IgE (*Ascaris*-sIgE), fecal examination for pathogenic intestinal helminths and tuberculin skin testing (TST). Histamine bronchoprovocation was tested in the group of children aged 10 years and older. Results were corrected for demographic variables, socioeconomic status, parental allergy, environmental tobacco smoke (ETS) exposure in the household, recent anthelminthic treatment and for clustering in the sampling unit.

Results Ascaris-sIgE was elevated in 48% of children, *Ascaris* eggs were found in 15% and TST was positive in 53%. Children with elevated *Ascaris*-sIgE had significantly increased risk of positive SPT to aeroallergens, particularly house dust mite, atopic asthma (*ever* and *recent*), atopic rhinitis (*ever* and *recent*) and increased atopy-related bronchial hyper-responsiveness. In children with negative TST (< 10 mm), elevated *Ascaris*-sIgE was associated with significantly increased risk of atopic symptoms (adjusted odds ratio (OR_{adj}) 6.5; 95% confidence interval (CI) 1.9–22.4), whereas in those with positive TST (\geq 10 mm) this association disappeared (OR_{adj} 0.96; 95% CI 0.4–2.8).

Conclusions These results suggest that immune response to *Ascaris* (*Ascaris*-sIgE) may be a risk factor of atopic disease in populations exposed to mild *Ascaris* infection and that MTB infection may be protective against this risk, probably by stimulation of anti-inflammatory networks.

Keywords *Ascaris* specific IgE, atopic disease, childhood, tuberculin skin test *Submitted 6 July 2005; revised 30 October 2005; accepted 17 January 2006*

Correspondence:

J.L.L. Kimpen, Department of Paediatrics, Wilhelmina Children's Hospital, University Medical Center, Lundlaan 6, PO Box 85090, 3508 AB Utrecht, The Netherlands. E-mail: ch.obihara@planet.nl

Introduction

Although infection with helminths can stimulate polyclonal immunoglobulin (Ig)E synthesis, their role in the development of atopic disease remains uncertain [1, 2]. The epidemiological relation of helminth infection and atopic disease, both associated with a T-helper (Th) 2 immune response, is controversial, as it has been reported that helminth infection may either suppress atopic disease [3–6] or predispose to it [7, 8]. Anthelminthic treatment may influence atopic conditions, or modulate the severity of symptoms or sensitization [9–11]. It is generally assumed that the prevalence of allergies is low in populations heavily infested with helminth [12–15].

The effect of intestinal helminth infection on atopic symptoms seems to depend on the duration and intensity

of infection [12, 13]. With chronic and intense infection, atopic symptoms may be suppressed, whereas mild and intermittent infection may result in an enhanced reaction to environmental allergens and atopic response [12, 13].

Previous epidemiological surveys often used different definitions of atopy, which made it difficult to compare results. Recently, the World Allergy Organization (WAO) published a new definition of atopic disease as clinical allergic symptoms in combination with a positive allergy skin prick test (SPT) and/or an elevated serum IgE antibody level [16], which makes it possible to define atopic diseases more uniformly.

Although *Mycobacterium tuberculosis* (MTB) infection is high in most helminth-infested populations in the developing world, the possible influence of MTB, a strong stimulator of Th1 immunity, on the effect of helminth infection on the development of atopic disease has not been investigated.

In a resource-poor urban suburb of South Africa, where helminth infection is endemic and tuberculosis (TB) incidence is high [17, 18], we investigated the associations of *Ascaris lumbricoides* infection (specific-IgE or eggs), and atopic sensitization to common aeroallergens or atopic diseases in the paediatric population, using the new WAO definitions. In addition, we tested the influence of a positive tuberculin skin test (TST) on the associations investigated.

Materials and methods

This cross-sectional study was conducted between June and October 2003, in an established epidemiological research site in Cape Town, South Africa, with a population of 36 334 in 2001 (Statistics South Africa: Western Cape, Census 2001) and predominantly of mixed ancestry. Although most of the families live in brick houses and have access to clean water and electricity, the general socioeconomic conditions are poor. The prevalence of childhood asthma is 10.8–13.3% and that of allergic rhinitis (AR) is 16% [19, 20]. The vaccination policy in South Africa requires all children to receive a single dose of *M. bovis* BCG-vaccine during the first week of life. On the basis of the World Health Organization (WHO) criteria, neonatal BCG vaccination in the study area is universal (\geq 90%).

All children aged 6–14 years living in a 15% randomly selected sample of household addresses were enumerated and invited to participate in the study. Exclusion criteria included known immune-compromising disease (such as HIV), active TB disease, pregnancy or parental refusal. Parents or legal guardians gave written, signed, informed consent.

Under the supervision of a trained field worker, each parent completed a written validated International Study of Asthma and Allergies in Children (ISAAC) Phase I questionnaire on previous (ever) and recent (< 12 months) symptoms of asthma or AR. In addition, the questionnaire included questions on socioeconomic variables, parental allergic history, BCG immunization, anthelminthic treatment, environmental tobacco smoke (ETS) exposure in the household and HIV status. The proportion of HIV-1-infected persons in the area is among the lowest in South Africa [21]. Children with questionnaire-reported positive HIV status were excluded, because of the possible effect on T cell immunity. The presence of a BCG scar was noted in each child.

SPT to eight common aeroallergens (ALK-Abelló, Hørsholm, Denmark) based on local allergen exposure (house dust mite (HDM), Bermuda grass, Rye grass, cat dander, dog dander, *Alternaria alternata*, *Cladosporium herbarium* and *Aspergillus*), together with a positive (histamine chloride 10 mg/mL) and negative (glycerol) control, were performed after completion of the questionnaire. A positive SPT reaction was defined as a mean weal diameter of \geq 3 mm in excess of the negative control.

Total serum IgE and *Ascaris*-sIgE levels were measured with the CAP RAST (Pharmacia[®], Uppsala, Sweden). Owing to the lack of reference values for non-Caucasian children for this analysis, total serum IgE level higher than the median value of the children in the study and *Ascaris*sIgE level ≥ 0.35 IU/mL were considered elevated. Two stool samples, taken at least 24 h apart, were collected from each child. A portion (0.5–1 g) was weighed, added to formalin, concentrated by formalin-ether sedimentation and examined by light microscopy. The presence of helminth eggs was defined as helminth infection. Helminth infection intensity (eggs per gram (epg) stool) was classified according to the proposed WHO classification (WHO/CTD/SIP/98.1).

The histamine bronchoprovocation test was used to assess bronchial hyper-responsiveness (BHR). The spirometry (with a calibrated Jaeger Masterscope, software version: 4.52i) was repeated until the best of three reproducible baseline measurements of forced expiratory volume at 1 s (FEV₁) was obtained; this was taken as the reference. During the pilot spirometry testing, a majority of the children less than 10 years of age failed to achieve reproducible readings within the allocated time frame of the test. This was mainly due to lack of motivation or difficulty with understanding the instructions. Because of this difficulty, the study coordinators decided to limit the histamine bronchoprovocation test to all children 10 years of age and older. The test was conducted according to the modified Cockcroft protocol, as described by Steinbrugger [22]. It consisted of a 2-minute inhalation of nebulized histamine (Pari LC PLUS nebulizer) through a mouthpiece of a Pari-Boy (37:00; 50 Hz; 1.3 bar2700 L/min; Pari, Starnberg, Germany). During inhalation, the nose was closed by a clip. The histamine concentration (0.03-7.8 mg/mL) was doubled in a standardized way, and the FEV₁

was measured after inhalation of each concentration. The test was continued until a decrease in FEV₁ from baseline of 20% or the maximum dose was reached. The response to the histamine challenge was expressed as PC_{20} ; the concentration that caused a fall in FEV₁ of 20% from baseline was calculated by linear interpolation of a log-linear dose-response curve. A subject was regarded as having an increased BHR if PC_{20} was $\leq 8.0 \text{ mg/mL}$. Children with an FEV₁ fall of $\geq 10\%$ after histamine challenge were treated with two inhalations of 200 µg salbutamol and FEV₁ measurement was repeated 5 min later.

The research site has a very high TB incidence [18]. The TB notification rate was 341 per 100 000 for new smearpositive TB and 612 per 100 000 for bacteriologicalconfirmed TB in 2002. Tuberculin reactivity becomes apparent in 3–6 weeks after initial MTB infection and may remain positive for the lifetime of the individual [23]. A positive TST reaction is an accepted hallmark of primary infection with MTB [23]. TST response was documented in the children. TST was performed by injecting 2 TU (tuberculin units) of PPD RT 23 (Statens Serum Institut, Copenhagen, Denmark) and measuring the transverse induration diameter after 2–3 days. In accordance with the American Thoracic Society guidelines, a positive TST was defined as an induration of \ge 10 mm [24].

Definition of atopic disease outcome variables

Allergic symptoms represent ISAAC-questionnaire-reported symptoms. A positive SPT was used to differentiate children with atopic symptoms from those with nonatopic symptoms. Atopic asthma and rhinitis were defined according to the new WAO definition, as questionnairereported asthma or AR together with a positive SPT [16]. Atopy-related BHR was defined as increased BHR with positive SPT (BHR with SPT).

Statistical analysis

Bivariate data analyses were performed using the χ^2 test (SPSS 11.0). Regression analyses were performed with the generalized estimated equation (GEE) logistic regression (LR) (STATA 8.0), using reported allergic symptoms, SPT reactivity, atopic diseases or BHR with SPT as the dependent variable and *Ascaris*-sIgE as the independent variable. The association was adjusted for possible confounding variables: demographic (age and gender), genetic (parental allergic history), socioeconomic – (household income), environmental (ETS exposure in the household) factors and anthelminthic treatment. In addition, the association was stratified for the presence or absence of a positive TST, in an LR model. Results were corrected for clustering (> 1 child per household) in the sampling unit. The sampling unit in the study is the

household. There were a total of 201 households in the study, with an average of $1 \cdot 5$ children per household. For the analysis, each child was coded and linked to the specific household. The study protocol was approved by the Ethics Review Board of the University of Stellenbosch.

Results

Of 418 enumerated children aged 6–14 years, 359 (86%) were enrolled in the study, 39 had moved away from the area, 14 refused consent, three were pregnant and three exceeded the age limit. The median age was 11.0 (range 6–14) years. The children excluded from the study did not differ in age or gender from those included.

Table 1 describes the most important characteristics of the 359 children analysed, together with prevalences of reported allergic symptoms, atopic sensitization, atopic diseases, increased BHR, helminth infection and anthelminthic treatment, serum IgE antibody levels, parental allergic history, positive TST reaction, ETS exposure in the household and BCG immunization. All children in the study reported to have received neonatal BCG

Table 1. Characteristics of the study population (total = 359)

	Number/total	0/0
Demographics		
Sex		
Male	188/359	52.4
Female	171/359	47.6
Age (years)		
6-10	167/359	46.5
11-14	192/359	53.5
Respiratory allergic symptoms		
Asthma		
Ever		
Yes	49/359	13.6
Recent (≤ 12 months)		
Yes	27/359	7.5
Allergic rhinitis		
Ever		
Yes	41/359	11.4
Recent		
Yes	34/359	9.5
Atopic sensitization:		
Positive SPT (\geq 3 mm)		
House dust mite	48/359	13.4
Rye grass	17/359	4.7
Alternaria	15/359	4.2
Bermuda grass	12/359	3.3
Cat	9/359	2.5
Aspergillus	4/359	1.1
Dog	4/359	1.1
Cladosporium	3/359	0.8
Any allergen	66/359	18.4
Respiratory atopic diseases*		

Table 1. Continued.

	Number/total	0/0
Atopic asthma		-
Ever		
Yes	16/359	4.5
Recent		
Yes	11/359	3.1
Atopic rhinitis		
Ever		
Yes	15/359	4.2
Recent		
Yes	14/359	3.9
$Histamine\ bronchoprovocation^{\dagger}$		
Increased BHR		
Yes	133/242	54.9
Increased BHR with positive SPT		
Yes	32/242	13.3
Helminth infection		
Helminth egg in stool sample		
Ascaris lumbricoides	53/359	14.8
Trichuris trichiura	144/359	40.1
Anthelminthic treatment $\leq 12 \text{ mon}$	ths	
Yes	75/359	20.9
Serum IgE antibody levels (IU/mL)		
Ascaris-specific-IgE (≥0.35)	171/357	47.9
Total IgE (≥median: 143.0)	178/357	49.9
Parental allergic history		
Yes	83/359	23.1
Environmental exposures		
Positive tuberculin skin test (TST)		
≥10 mm	179/337	53.1
Environmental tobacco smoke (ETS) exposure in the house	ehold
Yes	217/359	60.4
BCG immunization		
Yes	359/359	100

*Defined as combination of reported allergic symptom and a positive skin prick test (SPT), according to the new World Allergy Organization nomenclature.

[†]Histamine challenge was performed only in children \ge 10 years of age (explanation in the methods section).

BHR, bronchial hyper-responsiveness; IgE, immunoglobulin E.

immunization. The prevalence of BCG scar did not differ between children with positive TST and those with negative TST.

Of children with helminth infection, 14.8% (53) were infected with *Ascaris* and 40% (144) with *Trichuris trichiura (Trichuris)*. Other helminth eggs found were *Enterobius vermicularis* in four children and *Hymenolepsis nana* in one. The median infection intensity and IgE values were log-transformed because the values were not normally distributed. The median (range) infection intensity among the children infected with *Ascaris* was 182 (0–5888) epg and 58 (0–10 000) epg for *Trichuris*. The median (range) total IgE level was 143.00 IU/mL (2.33–18 080.00) and *Ascaris*-SIgE 0.35 IU/mL (< 0.35–63.10). Fifty percent (178) of children had an elevated

total IgE level and 48% (171) had an elevated *Ascaris*-sIgE level. Of children who underwent histamine bronchoprovocation, 55% (133) had increased BHR and 13.3% (32) had both increased BHR and positive SPT. TST was positive in 53% (179) of children. The TST distribution of the children was unimodal and the median TST size in those with any TST induration was 18 mm (range 1–30.5). The prevalence of parental allergic history or recent (< 12 months) anthelminthic treatment did not differ between children with elevated *Ascaris*-sIgE or *Ascaris* eggs in the stool and those without.

Children with *Ascaris* eggs in the stool were more likely to have an elevated *Ascaris*-sIgE level (adjusted odds ratio (OR_{adj}) 2.4; 95% confidence interval (CI) 1.3–4.5), and had a higher mean log *Ascaris*-sIgE (P < 0.005) than those without. In contrast, children with *Trichuris* infection were not more likely to have an elevated *Ascaris*-sIgE level (OR_{adj} 1.50; 95% CI 0.92–2.80) or a higher mean log *Ascaris*-sIgE (P = 0.10).

Ascaris-sIgE and atopic disease

Children with elevated Ascaris-sIgE had a higher frequency of allergic symptoms, atopic diseases, positive SPT to HDM, BHR and BHR with SPT than those without elevated Ascaris-sIgE. For reported asthma ever (P < 0.05), atopic asthma ever (P < 0.05) and recent (P < 0.005), atopic rhinitis ever (P < 0.05) and recent (P < 0.05), positive SPT to HDM (P < 0.05) and BHR (P < 0.05), the differences were significant. Figure 1 shows a significant linear association of log Ascaris-sIgE and the frequency of atopic diseases, positive SPT to HDM or BHR with SPT in the children. Table 2 shows that children with elevated Ascaris-sIgE had significantly increased risk of atopic sensitization, particularly HDM and atopic diseases, than those without. They also showed a trend to have a higher risk of reported allergic symptoms, BHR and BHR with SPT. The increase in risk of atopic disease or positive SPT to HDM observed in children with elevated Ascaris-sIgE remained significant after adjusting for possible confounding variables (Table 3). In a linear regression analysis, log Ascaris-sIgE was significantly inversely related to log PC20 (regression coefficient [β] = -0.206, SE 0.06, P = 0.02).

Children with *Ascaris* eggs in the stool did not have an increased risk of positive SPT (OR 0.57; 95% CI 0.23–1.40), atopic asthma (OR 0.85; 95% CI 0.81–0.86), atopic rhinitis (OR 1.04; 95% CI 0.22–4.82) or atopy-related BHR (OR 0.74; 95% CI 0.21–2.60) than those without. This did not change after adjusting for the possible confounding variables.

TST, Ascaris-sIgE and Atopic disease

Adjusted stratified analysis in an LR model, according to TST reactivity, showed that in children with negative TST



Fig. 1. Frequency of atopic diseases, positive skin prick test (SPT) to house dust mite and bronchial hyper-responsiveness with positive SPT showing direct linear increase with log *Ascaris* specific immunoglobulin (Ig)E (*Ascaris*-specific IgE) level (*P*-value for linear relation).

elevated *Ascaris*-sIgE was associated with increased risk of atopic symptoms (OR_{adj} 6.5; 95% CI 1.9–22.4), while in those with a positive TST elevated *Ascaris*-sIgE was not associated with increased risk (OR_{adj} 0.96; 95% CI 0.4–2.8). There was a significant interaction factor between a positive TST and an elevated *Ascaris*-sIgE (P = 0.02) in the LR model.

Discussion

This study in children from a resource-poor urban suburb of South Africa shows a significant association of elevated *Ascaris*-sIgE level and increased risk of atopic asthma, atopic rhinitis or hypersensitivity to common aeroallergens. It also shows that a positive TST, a hallmark of MTB infection, may influence the association of *Ascaris*-sIgE and atopic disease. In children with negative TST, elevated *Ascaris*-sIgE was associated with increased risk of atopy, whereas in those with positive TST no increased risk was observed.

These findings suggest that elevated *Ascaris*-sIgE may be a risk factor of respiratory atopic disease and is in agreement with previous studies in South Africa [8, 14, 25], South America [10] and China [7].

Although it has been hypothesized that polyclonal IgE produced during helminth infection may block allergic reaction by either suppressing antigen-specific IgE production or by saturation of IgE receptors on mast cells [10, 26, 27, 28], other studies that demonstrated lack of saturation of mast cell capacity *in vitro* or *in vivo* argue against this mechanism [29, 30]. Recent evidence suggests that chronic infections, such as helminth and MTB, may lead to CD4⁺ regulatory T cell (Treg) stimulation with subsequent production of high levels of anti-inflammatory cytokines, which may inhibit allergic inflammation [13]. This effect may depend on the intensity and persistence of infection [13, 31]. With intense and persistent

infection, atopic responses may be suppressed by strong stimulation of Tregs and anti-inflammatory cytokines [13]. In contrast, mild helminth infection (as in our study population) may lead to moderate or no stimulation of anti-inflammatory networks and this may result in an enhanced reaction to environmental allergens and atopic response [13, 31].

The mild intensity of *Ascaris* infection in our study population may be explained by the frequent single-dose anthelminthic therapy with benzimidazoles being used by children in the area [32]. This reduces the intensity of parasitic infestation and interrupts the persistence (or chronicity) of infection. The higher prevalence of *Trichuris* infection, which is less sensitive to a single dosage of anthelminthic therapy, supports this assumption [32].

A rather surprising finding is that the presence of Ascaris eggs in the stool was not associated with increased prevalence of atopic disease. This observation may reflect important differences in the immunological profile between Ascaris eggs in the stool and elevated Ascaris-sIgE, especially in populations where intestinal helminth infection is of mild intensity. This is supported by reports that an elevated Ascaris-sIgE level and the presence of Ascaris eggs in the stool may measure different entities [33]. An elevated Ascaris-sIgE level reflects the competency of the host Th2 immune response to Ascaris antigen. The presence of Ascaris eggs in the stool reflects active infection, although the impact of this infection on the host immune response may depend on different factors, including the severity and chronicity of infection [13, 31]. The lack of association between Ascaris eggs and atopic disease may partly be related to the low prevalence and mild intensity of Ascaris eggs in the children.

Our findings differ from those of another research group, who reported a lower prevalence of hyper-

Table 2. Crude and adjusted risk of allergic symptoms, atopic sensitization and atopic diseases in all children and in those *with* elevated *Ascaris*-specific IgE (*Ascaris*-sIgE ≥ 0.35 IU/mL)

		Children with elevated				
	All children ($n = 359$)	Ascaris-sIgE ($n = 171$)	Crude OR	95% CI	Adjusted OR*	95% CI
Allergic symp	otoms and BHR					
Asthma						
Ever						
No	308	141 (45.8)	1		1	
Yes	49	30 (61.2)	1.87	1.01-3.46	1.79	0.90-3.53
Recent						
No	330	155 (47.0)	1		1	
Yes	27	16 (59.3)	1.64	0.74-3.65	1.55	0.64-3.77
Allergic rhini	tis					
Ever						
No	316	147 (46.5)	1		1	
Yes	41	24 (58.5)	1.62	0.84-3.13	1.51	0.71-3.18
Recent						
No	323	150 (46.4)	1		1	
Yes	34	21 (61.8)	1.86	0.90-3.85	1.9	0.85-4.26
Increased BH	R					
No	109	41 (37.6)	1		1	
Yes	131	67 (51.1)	1.74	1.04-2.91	1.67	0.93-3.00
Atopic sensit	ization					
HDM						
No	309	138 (44.7)	1		1	
Yes	48	33 (68.8)	2.73	1.42-5.22	3.81	1.76-8.26
Any positive	SPT					
No	292	133 (45.5)	1		1	
Yes	65	38 (58.5)	1.68	0.98-2.90	2.09	1.15-3.79
Atopic diseas	es [†]					
Atopic asthm	a					
Ever						
No	341	159 (46.6)	1		1	
Yes	16	12 (75.0)	3.4	1.1-10.9	3.97	1.19-13.23
Recent						
No	346	161 (46.5)	1		1	
Yes	11	10 (90.9)	11.5	1.5-90.7	14.26	1.78-114.02
Atopic rhiniti	is					
Ever						
No	343	160 (46.8)	1		1	
Yes	15	11 (73.3)	3.1	1.0-10.0	4.56	1.19-17.48
Recent						
No	342	160 (46.6)	1		1	
Yes	14	11 (78.6)	4.2	1.2-15.3	8.64	1.91-39.11
Increased BH	R with SPT [‡]					
No	208	89 (42.8)	1		1	
Yes	32	19 (59.4)	2.0	0.9-4.2	2.08	0.90-4.84

*Adjusted odds ratios (OR) were calculated with the logistic regression model (generalized estimated equation, GEE) using allergic symptoms, atopic sensitization, atopic diseases or BHR+SPT as dependent variables and elevated *Ascaris*-sIgE together with confounding variables: age, gender, parental allergic history, average household income, antihelminthic treatment in the past 12 months, environmental tobacco smoke (ETS) exposure in the household and tuberculin skin test reactivity as independent variables. In addition, correction was made for clustering (> 1 child per household) in the sampling unit.

[†]Atopic disease was defined using the new World Allergy Organization nomenclature as reported allergic symptom and a positive skin prick test (SPT) to common aeroallergens.

[‡]Only 240 children aged \ge 10 years of age with complete data on histamine bronchoprovocation and *Ascaris*-sIgE were included in the analysis. BHR, bronchial hyper-responsiveness; IgE, immunoglobulin E; HDM, house dust mite.

	Adjusted OR (95% CI)						
Variable (categories)	Atopic asthma ever	Recent atopic asthma	Atopic rhinitis ever	Recent atopic rhinitis	Positive SPT to HDM		
Ascaris-sIgE (IU/mL)							
< 0.35	1.0	1.0	1.0	1.0	1.0		
≥0.35	3.97 (1.19–13.23)	14.26 (1.78-114.02)	4.56 (1.19-17.48)	8.64 (1.91-39.12)	3.81 (1.76-8.26)		
Age (years)							
6-10	1.0	1.0	1.0	1.0	1.0		
11-14	1.83 (0.55-6.13)	3.0 (0.60-14.89)	3.04 (0.71-12.94)	4.94 (0.98-25.04)	2.26 (1.04-4.89)		
Gender							
Male	1.0	1.0	1.0	1.0	1.0		
Female	1.75 (0.59-5.21)	1.17 (0.86-15.06)	3.61 (0.07-1.16)	2.99 (0.70-12.73)	1.85 (0.89-3.85)		
Household income (SA	R)						
< 800	1.0	1.0	1.0	1.0	1.0		
≥800	2.04 (0.48-8.62)	2.72 (0.52-14.16)	3.99 (0.94-16.88)	6.16 (1.17-32.43)	1.84 (0.65-5.20)		
Parental allergy							
No	1.0	1.0	1.0	1.0	1.0		
Yes	1.75 (0.55–5.56)	1.29 (0.30-5.57)	1.13 (0.26-4.91)	1.56 (0.33-7.40)	1.46 (0.65-3.30)		
TST (mm)							
< 10	1.0	1.0	1.0	1.0	1.0		
≥10	0.62 (0.21-1.83)	0.43 (0.12-1.58)	0.12 (0.02-0.64)	0.05 (0.01-0.41)	0.89 (0.43-1.82)		
Recent helminthic treat	tment						
No	1.0	1.0	1.0	1.0	1.0		
Yes	0.29 (0.03–2.50)	0.48 (0.05-4.67)	0.71 (0.13-3.96)	0.98 (0.16-5.98)	1.18 (0.48-2.93)		
Current smoke exposur	re						
No	1.0	1.0	1.0	1.0	1.0		
Yes	0.92 (0.30–2.88)	0.81 (0.21–3.10)	0.24 (0.07–0.90)	0.17 (0.04–0.79)	0.26 (0.26-1.20)		

Table 3. Adjusted risk of atopic disease outcomes, positive skin test to house dust mite (HDM) and *Ascaris*-specific IgE (*Ascaris*-sIgE), together with confounding variables used in the logistic regression model

Adjusted odds ratios (OR_{adj}) were calculated with the generalized estimated equation (GEE) logistic regression (LR) model (STATA 8.0), using atopic asthma, atopic rhinitis and positive SPT to HDM as the dependent variables and *Ascaris*-sIgE and the other confounding variables in the table as independent variables. In addition, correction was made for clustering (> 1 child per household) in the sampling unit. Atopic disease was defined using the new World Allergy Organization nomenclature as reported allergic symptom and a positive skin prick test (SPT). CI, confidence interval; IgE, immunoglobulin; tuberculin skin testing.

sensitivity to HDM in African children with urinary schistosomiasis [30]. The authors considered the antiinflammatory cytokine, IL-10, produced during *Schistosoma* infection to be responsible for the effect. The enhanced production of IL-10 seen with chronic tissue helminth infections such as filariasis and schistosomiasis has not been observed in ascariasis [34]. This, and the mild intensity of helminth infection, may account for the different observations.

Consistent with previous findings, we found that a positive TST was inversely associated with the risk of atopic disease [35]. This is in support of recent evidence that chronic infections, such as MTB, may down-regulate the atopic response through the stimulation of anti-inflammatory networks [13, 31]. The strong median TST reaction (18 mm) in the children indicates hypersensitivity to natural MTB infection that is highly prevalent in the community and not to BCG or environmental mycobacteria, which generally stimulate a weaker TST reaction [23, 24, 35, 36]. Our present findings contrast with the results of a study in Gambian children, in which no inverse

association was found between positive TST and atopy in children [37]. The difference in findings could be explained by the fact that a larger proportion of children in our study had a strongly positive TST size than in the Gambian study. This is supported by observations that the size of TST is directly related to serum IFN- γ level, suggesting that increased serum IFN- γ concentration may partly explain the inverse association between MTB infection and atopy [38]. Furthermore, the new globally accepted WAO definition of atopic rhinitis was used in our study [16]. This contrasts with the Gambian study in which atopy was defined as a positive SPT only.

A particular strength of this study is the use of objective measures (SPT) to differentiate children with atopic and non-atopic symptoms, based on the new WAO definitions [16]. Most previous studies defined atopic outcome either as a positive SPT or an elevated serum IgE only [6, 11]. The influence of MTB infection was taken into account, and this, to our knowledge, has not been reported previously in this context. The main limitations are the cross-sectional design and questionnaire-based diagnosis of allergic symptoms. Limitations inherent to the crosssectional design are recall-bias and the inability to analyse the temporal relationship of Ascaris eggs, AscarissIgE, the development of atopic disease and TST reactivity. Questionnaire-based limitations were reduced by using the validated ISAAC questionnaire and collecting questionnaire data before performing objective tests. The reliability of questionnaire-generated data is supported by the fact that the prevalences of allergic symptoms were comparable with those of the ISAAC study in Cape Town [20]. That reliable spirometric tests could not be achieved in younger children does not bias the results, but rather reflects an important difficulty that may be encountered in performing such a standardized test in otherwise healthy voluntary children. Although the factors responsible for the high frequency of increased BHR were not investigated, it may be related to the high prevalence of ETS exposure and other non-atopic environmental airway pollutants in the area. ETS is a known risk factor for airway hyper-responsiveness in children [39].

In conclusion, this study shows a significant association of elevated *Ascaris*-sIgE and atopic sensitization and symptoms, suggesting that elevated *Ascaris*-sIgE may be an environmental risk factor of atopic manifestation, particularly in helminth-infected children with mild infection. It also shows the influence of natural MTB infection on the association, which may be attributed to stimulation of anti-inflammatory immune networks during MTB infection. There is a need for prospective community-based studies to confirm these findings.

Acknowledgements

We thank all the parents and children who participated in this study. We are indebted to Prof. P.C. Potter (Allergy Diagnostic and Clinical Research Unit, University of Cape Town, South Africa) for the serum IgE analyses and allergy diagnostics. We are also indebted to Ann Toerien for carrying out the allergy and tuberculin skin tests, Jerome Cornelius for the successful fieldwork and Dr Vera Adams (Nutritional Intervention Research Unit, Medical Research Council of South Africa, Cape Town, South Africa). We also thank Edwin Videler and Johan Mouton (Lung Function Laboratory of the Department of Respiratory Medicine, Stellenbosch University), and Rita van Deventer (Parasitology Reference Unit, National Institute for Communicable Diseases, National Health Laboratory Services, Johannesburg, South Africa).

Sources of support and grants: C.C.O is a recipient of a grant from the Ter Meulen Fund, Royal Netherlands Academy of Arts and Sciences. This study was funded by the Stellenbosch University (through funding from the South African Department of Trade and Industry, THRIP fund) and the Wilhelmina Children's Hospital, University Medical Center Utrecht, the Netherlands.

References

- 1 Britton J. Parasites, allergy, and asthma. *Am J Respir Crit Care Med* 2003; 168:266–7.
- 2 King C, Low C, Nutman T. IgE production in human helminth infection. Reciprocal interrelationship between IL-4 and IFNgamma. *J Immunol* 1993; 150:1873–80.
- 3 Cooper PJ, Chico ME, Bland M, Griffin G, Nutman B. Allergic symptoms, atopy, and geohelminth infections in a rural area of Ecuador. Am J Respir Crit Care Med 2003; 168:313–7.
- 4 Nyan OA, Walraven G, Banya W *et al*. Atopy, intestinal helminth infection and total serum IgE in rural and urban adult Gambian communities. *Clin Exp Allergy* 2001; 31:1672–8.
- 5 Scrivener S, Yemaneberhan H, Zebenigus M *et al.* Independent effects of intestinal parasite infection and domestic allergen exposure on the risk of wheeze in Ethiopia: a nested case–control study. *Lancet* 2001; 358:1493–9.
- 6 Dagoye D, Bekele Z, Woldemichael K *et al.* Wheezing, allergy, and parasite infection in children in urban and rural Ethiopia. *Am J Respir Crit Care Med* 2003; **167**:1369–73.
- 7 Palmer LJ, Celedón J, Weiss ST, Wang B, Fang Z, Xu X. *Ascaris lumbricoides* infection is associated with increased risk of child-hood asthma and atopy in rural China. *Am J Respir Crit Care Med* 2002; **165**:1489–93.
- 8 Joubert JR, De Klerk HC, Malan C. *Ascaris lumbricoides* and allergic asthma: a new perspective. *S Afr Med J* 1979; **56**:599–602.
- 9 Lynch NR, Palenque M, Hagel I, DiPrisco M. Clinical improvement of asthma after anthelminthic treatment in a tropical situation. *Am J Respir Crit Care Med* 1997; 156:50–4.
- 10 Lynch NR, Hagel I, Perez M, Di Prisco M, Lopez R, Alvarez N. Effect of anthelmintic treatment on the allergic reactivity of children in a tropical slum. *J Allergy Clin Immunol* 1993; 92:404–11.
- 11 Van den Biggelaar AHJ, Rodrigues LC, van Ree R et al. Long-term treatment of intestinal helminths increases mite skin-test reactivity in Gabonese schoolchildren. J Infect Dis 2004; 189:892–900.
- 12 Cooper PJ. Can intestinal helminth infections (geohelminths) affect the development and expression of asthma and allergic disease. *Clin Exp Immunol* 2002; **128**:398–404.
- 13 Yazdanbakhsh M, Van den Biggelaar, Maizels RM. Th2 responses without atopy: immunoregulation in chronic helminth infections and reduced allergic disease. *Trends Immunol* 2001; 22:372–7.
- 14 Joubert JR, van Schalkwyk DJ, Turner KJ. *Ascaris lumbricoides* and human immunogenic response: enhanced IgE-mediated reactivity to common inhaled allergens. *S Afr Med J* 1980; 57:409–12.
- 15 Lynch NR, Medouze L, di Prisco-Fuenmayor MC, Verde O, Lopez RI, Malave C. Incidence of atopic disease in a tropical environment: partial independence from intestinal helminthiasis. *J Allergy Clin Immunol* 1984; 73:229–33.
- 16 Johansson SGO, Bieber T, Dahl R *et al.* Revised nomenclature for allergy for global use: report of the nomenclature review committee of the world allergy organization, October 2003. *J Allergy Clin Immunol* 2004; 113:832–6.
- 17 Gunders AE, Cotton M, Nel E *et al*. Prevalence and intensity of intestinal worm infections in crèche attenders in urban and periurban settings in greater Cape Town. *S Afr J Epidemiol Infect* 1993; 8:48–51.

- 18 Verver S, Warren RM, Munch Z. *et al.* Proportion of tuberculosis transmission that takes place in households in a high-incidence area. *Lancet* 2004; **363**:212–4.
- 19 Ehrlich RI, Du Toit D, Jordaan E, Volmink JA, Weinberg EG, Zwarenstein M. Prevalence and reliability of asthma symptoms in primary school children in cape town. *Int J Epidemiol* 1995; 24:1138–45.
- 20 The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema. *Lancet* 1998; 351:225–32.
- 21 National HIV and Syphilis antenatal sero-prevalence survey in South Africa 2002. The South African Department of Health. http://www.doh.gov.za/docs/reports/2002/hiv-syphilis.pdf
- 22 Steinbrugger B, Eber E, Modl M, Weinhandl E, Zach S. A comparison of a single-step cold air challenge and routine histamine provocation for the assessment of bronchial hyperresponsiveness in children and adolescents. *Chest* 1995; 103:741–5.
- 23 Shingadia D, Novelli V. Diagnosis and treatment of tuberculosis in children. *Lancet Infect Dis* 2003; 3:624–32.
- 24 American Academy of Paediatrics. Pickering LK, editor *Red Book: 2003 Report of the Committee on Infectious Diseases*, 26th Edn. Elk Grove Village, IL: American Academy of Paediatrics, 2003; 643.
- 25 Joubert JR, Brink S, Hentzen GM. Allergic asthma in different population groups in the Western Cape. *S Afr Med J* 1988; 73:150–4.
- 26 Mitre E, Norwood S, Nutman TB. Saturation of immunoglobulin E (IgE) binding sites by polyclonal IgE does not explain the protective effect of helminth infections against atopy. *Infect Immun* 2005; **73**:4106–11.
- 27 Godfrey RC, Gradidge CF. Allergic sensitization of human lung fragments prevented by saturation of IgE binding sites. *Nature* 1976; 259:484–6.
- 28 Hagel I, Lynch NR, DiPrisco MC, Lopez R, Garcia N. Allergic reactivity of children of different socio-economic levels in tropical populations. *Int Arch Allergy Immunol* 1993; 101:209–14.

- 29 Larrick JW, Buckley CF, Machamer CE *et al.* Does hyperimmunoglobulinemia-E protect tropical populations from allergic disease. *J Allergy Clin Immunol* 1983; 71:184–8.
- 30 Van den Biggelaar AH, van Ree R, Rodriguez LC et al. Decreased atopy in children infected with Schistosoma haematobium: role for parasite-induced interleukin-10. Lancet 2000; 356:1723-7.
- 31 Matricardi PM, Yazdanbakhsh M. Mycobacteria and atopy, 6 years later: a fascinating, still unfinished, business. *Clin Exp Allergy* 2003; 33:717–20.
- 32 Fincham JE, Markus MB, Adams VJ *et al.* Association of deworming with reduced eosinophilia: implications for HIV/AIDS and co-endemic diseases. *S Afr J Sci* 2003; 99: 182–3.
- 33 Dold S, Heinrich J, Wichmann H-E, Wjst M. Ascaris-specific IgE and allergic sensitization in a cohort of school children in the former East Germany. J Allergy Clin Immunol 1998; 102:414–20.
- 34 Cooper PJ, Chico ME, Sandoval C *et al.* Human infection with *Ascaris lumbricoides* is associated with a polarized cytokine response. *J Infect Dis* 2000; **182**:1207–13.
- 35 Shirakawa T, Enomoto T, Shimazu S, Hopkin JM. The inverse association between tuberculin responses and atopic disorder. *Science* 1997; 275:77–9.
- 36 Obihara CC, Beyers N, Gie RP *et al.* Inverse association between Mycobacterium tuberculosis infection and atopic rhinitis in children. *Allergy* 2005; **60**:1121–5.
- 37 Ota M, van der Sande M, Walraven G *et al.* Absence of association between delayed type hypersensitivity to tuberculin and atopy in children in The Gambia. *Clin Exp Allergy* 2003; 33:731–6.
- 38 Black GF, Fine PEM, Warndorff DK *et al.* Relationship between IFN-γ nd skin test responsiveness to mycobacterium tuberculosis PPD in healthy, non-BCG-vaccinated young adults in Northern Malawi. *Int J Tuberc Lung Dis* 2001; 7:664–72.
- 39 Stocks J, Dezateux The effect of parental smoking on lung function and development of the child. *Respirology* 2003; 8:266–85.

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.