

SAMRC InfoSpace

Acceptability of self- collection for human papillomavirus detection in the Eastern Cape, South Africa

Item Type	Article
Authors	Taku, O.;Meiring, T.L.;Gustavsson, I.;Phohlo, K.;Garcia-Jardon, M.;Mbulawa, Z.Z.A.;Businge, C.B.;Gyllensten, U.;Williamson, A.L.
Citation	Taku O, Meiring TL, Gustavsson I, Phohlo K, Garcia-Jardon M, Mbulawa ZZA, Businge CB, Gyllensten U, Williamson AL. Acceptability of self- collection for human papillomavirus detection in the Eastern Cape, South Africa. PLoS One. 2020 Nov 10;15(11):e0241781. doi: 10.1371/journal.pone.0241781.
DOI	10.1371/journal.pone.0241781
Publisher	PLoS One
Journal	PLoS One
Rights	Attribution 3.0 United States
Download date	2026-06-06 01:05:06
Item License	http://creativecommons.org/licenses/by/3.0/us/
Link to Item	https://pubmed.ncbi.nlm.nih.gov/33170891/

RESEARCH ARTICLE

Acceptability of self- collection for human papillomavirus detection in the Eastern Cape, South Africa

Ongeziwe Taku^{1,2}, Tracy L. Meiring^{1,2}, Inger Gustavsson³, Keletso Phohlo^{1,2}, Mirta Garcia-Jardon⁴, Zizipho Z. A. Mbulawa^{1,5,6,7}, Charles B. Businge^{8,9}, Ulf Gyllensten³, Anna-Lise Williamson^{1,2,5*}

1 Division of Medical Virology, Department of Pathology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa, **2** Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town, South Africa, **3** Department of Immunology, Genetics, and Pathology, Biomedical Center, Science for Life Laboratory Uppsala, Uppsala University, Uppsala, Sweden, **4** Department of Pathology at Walter Sisulu University and National Health Laboratory Service, Mthatha, South Africa, **5** SAMRC Gynaecological Cancer Research Centre, University of Cape Town, Cape Town, South Africa, **6** Department of Laboratory Medicine and Pathology, National Health Laboratory Service Nelson Mandela Academic Hospital and Walter Sisulu University, Mthatha, Eastern Cape, South Africa, **7** Medical Virology, National Health Laboratory Service, Nelson Mandela Academic Hospital, Mthatha, South Africa, **8** Department of Obstetrics and Gynaecology, Nelson Mandela Academic Hospital, Mthatha, South Africa, **9** Department of Obstetrics and Gynaecology, Faculty of Health Sciences, Walter Sisulu University, Mthatha, South Africa

* anna-lise.williamson@uct.ac.za



OPEN ACCESS

Citation: Taku O, Meiring TL, Gustavsson I, Phohlo K, Garcia-Jardon M, Mbulawa ZZA, et al. (2020) Acceptability of self- collection for human papillomavirus detection in the Eastern Cape, South Africa. PLoS ONE 15(11): e0241781. <https://doi.org/10.1371/journal.pone.0241781>

Editor: Marcia Edilaine Lopes Consolaro, Universidade Estadual de Maringa, BRAZIL

Received: July 2, 2020

Accepted: October 20, 2020

Published: November 10, 2020

Copyright: © 2020 Taku et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are in the [Supporting information](#) files.

Funding: This study was supported in part by the National Research Foundation of South Africa in the form of a grant awarded to ALW (64815), SAMRC-FORTE Collaborative Research Programme – South African Medical Research Council – Swedish Research Council for Health, Working Life and Welfare in the form of a grant awarded to ALW, and Poliomyelitis Research Foundation, South Africa in the form of bursaries

Abstract

Human papillomavirus (HPV) testing on vaginal self-collected and cervical clinician-collected specimens shows comparable performance. Self-sampling on FTA cards is suitable for women residing in rural settings or not attending regular screening and increases participation rate in the cervical cancer screening programme. We aimed to investigate and compare high-risk (HR)-HPV prevalence in clinician-collected and self-collected genital specimens as well as two different HPV tests on the clinician collected samples. A total of 737 women were recruited from two sites, a community health clinic (n = 413) and a referral clinic (n = 324) in the Eastern Cape Province. Cervical clinician-collected (FTA cards and Digene transport medium) and vaginal self-collected specimens were tested for HR-HPV using the *hpVIR* assay (FTA cards) and Hybrid Capture-2 (Digene transport medium). There was no significant difference in HR-HPV positivity between clinician-collected and self-collected specimens among women from the community-based clinic (26.4% vs 27.9%, $p = 0.601$) or the referral clinic (83.6% vs 79.9%, $p = 0.222$). HPV16, HPV35, and HPV33/52/58 group were the most frequently detected genotypes at both study sites. Self-sampling for HPV testing received a high positive response of acceptance (77.2% in the community-based clinic and 83.0% in referral clinic). The overall agreement between *hpVIR* assay and HC-2 was 87.7% ($k = 0.754$). The study found good agreement between clinician-collected and self-collected genital specimens. Self-collection can have a positive impact on a cervical screening program in South Africa by increasing coverage of women in rural areas, in

awarded to OT and KP. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Anna-Lise Williamson's salary comes from National Research Foundation of South Africa (Grant Number: 64815) The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: CC, cervical cancer; CI, confidence intervals, ref: reference; CIN, cervical intraepithelial neoplasia; HR-HPV, high-risk human papillomavirus; NILM, negative for intraepithelial lesions or malignancy; VIA, Visual Inspection Acetic acid.

particular those unable to visit the clinics and women attending clinics where cytology-based programs are not functioning effectively.

Introduction

Cervical cancer (CC), is the third-ranking cause of cancer disease with an estimated 569,847 cases leading to 311,365 deaths in women globally in 2018 [1]. The age-standardized incidence rate of cervical cancer varies worldwide from ≤ 2 to 75 per 100,000 women in different populations [2]. In South Africa, CC is the leading cause of cancer in women of reproductive age (15–44 years) and the most important cancer in black women. About 12,983 new cervical cancer cases are diagnosed annually in South Africa (estimates for 2018) with an incidence of 31.7 per 100,000 women [3, 4].

CC is causally associated with infection by high-risk types of human papillomavirus (HPV) [1]. Cervarix, which targets HPV16 and 18 [5], is the current HPV vaccine used in school-based HPV vaccination programme in South Africa. Although this vaccine is expected to prevent at least 70% of CCs, both vaccinated women and unvaccinated women still need to continue to be screened for CC [5]. CC screening methods include Visual Inspection with Acetic acid (VIA), cytology-based (Pap smear or liquid-based cytology) and HPV DNA testing [6]. Worldwide, cytology-based screening is still the most used screening test, although it has not been successfully implemented in African countries, including South Africa [7, 8]. In the Eastern Cape province of South Africa CC screening coverage is $< 50\%$, which is below the national target of 70% coverage [9]. Pap smear screening is not readily available to most of the women in rural South Africa, including Eastern Cape Province [9, 10]. In addition, while CC screening programmes may be available in some local clinics, the majority of women are not aware of the available services and not likely to participate [11–13].

HPV DNA testing is an effective CC prevention screening method and offers many advantages compared to cytology-based screening. This includes high sensitivity in predicting future precancerous lesions and CC, ability to perform self-collection of specimen for HPV testing, and the possibility to increase of screening intervals for women with a HPV negative test (> 5 -years) [14, 15]. A randomized study in India reported that a single HPV DNA test is more likely to lower the mortality rate of CC disease as compared to cytology-based screening or VIA [16]. Australia and Netherlands have introduced vaginal self-sampling for HPV DNA testing in their routine screening program as an alternative screening method for non-attendees [17, 18], and shown increased participation [19–21]. Therefore, introducing a self-collection method might overcome barriers that limit the use of the Pap smear test and make CC screening accessible to more women.

Liquid-based transport medium is commonly used to transport and store cervical or vaginal specimens for HPV testing. FTA cards for storage of cervical or vaginal specimens, have several advantages over a liquid-based media. These include inclusion of a color indicator to confirm that the sample is applied correctly, the samples become non-infectious once dried, long term stable storage at room temperature and easy transport [22–25]. Unlike the liquid-based transport medium, FTA cards are suitable for self-collection at home and can be transported via regular mail or other means not involving a cold-chain [22]. The process of extracting DNA from the FTA card is easy, quick and can be automated [22, 24]. Using the FTA card in combination with a clinically validated HPV DNA test has also been shown to work effectively for detection of HR-HPV infection [26]; with clinical specimens depicting up to 100% agreement of HR-HPV detection between FTA cards and liquid-based medium [27].

Due to the lack of infrastructure to perform Pap smear tests, cytology screening is not fully functional in rural settings. It is essential to implement a CC screening test that is easily accessible (such as self-collection) and simple to handle for women that do not have access to healthcare facilities offering regular cytology screening. In this study we aim to a) determine the prevalence of HR-HPV genotypes in women attending a community-based clinic and a referral clinic in the Eastern Cape, b) investigate and compare the HR-HPV prevalence in clinician-collected and self-collected genital specimens on FTA cards and to c) investigate attitudes to self-collection in women from the Eastern Cape.

Materials and methods

Study population and sample collection

This cross-section study was carried out from September 2017 to March 2019 at a community health clinic and the referral clinic for women with abnormal Pap smears within the OR Tambo district in the Eastern Cape Province of South Africa. A total of 413 women age ≥ 30 years attending the community health clinic for CC screening or others reasons were recruited. Also, 324 women age ≥ 18 years with abnormal cervical cytology and CC were recruited from the referral clinic. Written signed consent forms were obtained from all study participants, and sociodemographic data with other factors were obtained through a questionnaire. The ethical approval for this study was granted by the Human Research Ethics Committees of the University of Cape Town (UCT) (HREC reference 615/2017), Walter Sisulu University (reference 090/2016), and Eastern Cape Department of Health Ethics (EC reference 2017RP0_484).

Study participants provided three genital specimens. One clinician-collected cervical and one self-collected vaginal specimen were collected using Viba-brushes (Rovers Viba-Brush, Rovers Medical Devices) and applied onto indicating FTA elute cards (GE Health Care Life Sciences, Buckinghamshire, United Kingdom). A second clinician-collected cervical specimen was obtained using cervical brushes and stored in Digene transport medium (Qiagen, Gaithersburg, MD, USA). All specimens were kept at room temperature until shipped to the UCT HPV laboratory for analysis. Among participants recruited from the referral clinic, a cervical biopsy was collected and sent to the National Health Laboratory Service for histological analysis. The histology results were interpreted according to the guidelines of the International Agency for Research on Cancer [28].

Elution of DNA from FTA cards

DNA elution was performed as previously described [22]. Briefly, four 3.2 mm punches were obtained from each FTA card using the DBS Puncher Instrument (Perkin Elmer Life and Analytical Sciences, Wallac, Oy, Finland) and collected in a 96-well plate. The punches were washed with sterile water and the DNA eluted in 70 μ l water by incubating in a PCR machine at 95°C for 30 minutes. The DNA was stored at -20°C until further use.

Identification of HR-HPV using *hpVIR* real-time PCR

The eluted DNA from specimens on FTA cards was tested for HR-HPV using the clinically validated *hpVIR* real-time PCR assay [29]. *hpVIR* assay detects 12 HR-HPV genotypes (HPV16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58 and 59) and the analysis was performed as previously described [30]. Some HR-HPV types (16, -31, -35, -39, -51, -56, and -59) are identified as individual types, while HPV18 and -45, and HPV33, -52 and -58 are detected as two groups. The *hpVIR* assay also detect a human single copy house-keeping gene encoding Homo sapiens hydroxymethylbilane synthase (HMBS; GenBank accession no.M9523.1). This serves as a

control for the integrity of the eluted DNA and interpretation of the results for HPV-negative samples. The *hpVIR* assay has a cut-off of for HMBS single copy gene, and 10 copies per PCR of HPV for a positive HR-HPV type [29].

Detection of HR-HPV using Hybrid Capture-2

Cervical specimens in Digene transport medium were tested for 13 HR-HPV types (HPV16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59 and 68) using the Hybrid Capture-2 (HC-2) assay (Qia-gen Inc., Gaithersburg, MD; USA) according to the manufacturers protocol. A ratio of relative light units/cut-off ≥ 1 was considered positive while a ratio < 1 was considered negative for HR-HPV types.

Statistical analysis

Data analysis was performed using GraphPad Prism version 6 for Windows (Graphpad Software, La Jolla California USA). Descriptive statistics (medians and interquartile range) and frequency distribution were used to describe the sociodemographic and other variables of study participants in the population. For statistical analysis, chi-squared test was used to determine the difference in estimated HR-HPV prevalence between self-collected and clinician-collected samples. A p -value < 0.05 was considered significant. The overall agreement and HR-HPV genotype distribution agreement (percent agreement, kappa values with 95% Confidence Intervals) were determined using the kappa statistic. The kappa values were interpreted using a standard method (<http://www.graphpad.com/quickcalcs/Kappa2.cfm>).

Results

Description of study participants

Table 1 summarises the baseline demographics of study participants and other factors. Women from the community-based clinic had a median age of 46 years (IQR: 38–55), most were ≥ 50 years (42.4%), 37.3% were HIV-positive and the majority had been pregnant (96.6%). Almost half the women had ≥ 3 -lifetime sexual partners (49.4%) and most of the women never smoked (94.4%). A high proportion of women had attended high school or university (73.6%), and the monthly household income was $< \$139,36$ (71.2%).

Median age of the referral clinic women was 40.5 years (IQR: 33–49) and the majority were HIV-positive (71.3%). Half of the study participants had attended high school or university (54.3%) and 77.4% of women had a monthly household income of $< \$139,36$. The majority of women had their first sexual experience at the age of 16–18 years (54.3%), with 60.8% of them having ≥ 3 -lifetime sexual partners.

HR-HPV prevalence and agreement between self-collected and clinician-collected specimen using the *hpVIR* assay

All specimens on FTA cards had sufficient amounts of genomic DNA (10 copies or more of the HMBS single copy gene) for the HPV test to be informative. For women recruited at the community-based clinic, there was no significant difference in HR-HPV prevalence between self-collected and clinician-collected samples [27.9%(115/413) vs 26.4%(109/413), $p = 0.639$]. HR-HPV positivity between self-collected and clinician-collected samples showed an agreement of 86.9%, with a kappa value of 0.669 [95%CI: 0.588–0.750, (Table 2)].

Among the referral clinic women, HR-HPV prevalence was slightly higher in clinician-collected versus self-collected specimens, but the difference was not significant [83.6% (271/324) vs 79.9% (259/324), $p = 0.222$]. The observed overall agreement of HR-HPV infection between

Table 1. Demographic and behaviour characteristics of study participants.

	Community clinic	Referral clinic
Variables	% (n/N)	% (n/N)
HIV Status		
No	62.7 (259/413)	27.8 (90/324)
Yes	37.3 (154/413)	71.3 (231/324)
Missing	0.0 (0/413)	0.9 (3/324)
If yes, ARV's?		
No	3.9 (6/154)	1.3 (3/231)
Yes	96.1 (148/154)	98.3 (227/231)
Missing	0.0 (0/154)	0.4 (1/231)
Age categories		
18–29 years	13.0 (42/324)
30–39 years	31.5 (130/413)	33.3 (108/324)
40–49 years	26.2 (108/413)	31.2 (101/324)
≥50 years	42.4 (175/413)	22.5 (73/324)
Missing	0.0 (0/413)	0.0 (0/324)
Highest level of education attained		
Never/ primary	26.4 (109/413)	45.7 (148/324)
High school/university	73.6 (304/413)	54.3 (176/324)
Missing	0.0 (0/413)	0.0 (0/324)
Household income		
< \$139,36	71.2 (294/413)	77.5 (251/324)
≥ \$139,36	27.6 (114/413)	21.3 (69/324)
Missing	1.2 (5/413)	1.2 (4/324)
Smoking status		
Never	94.4 (390/413)	92.0 (298/324)
Former/current smoker	5.6 (23/413)	7.7 (25/324)
Missing	0.0 (0/413)	0.3 (1/324)
Ever drank alcohol		
No	91.0 (376/413)	81.5 (264/324)
Yes	9.0 (37/413)	18.2 (59/324)
Missing	0.0 (0/413)	0.3 (1/324)
Age at first sexual experience		
<16 years	12.1 (50/413)	19.4 (63/324)
16–18 years	52.5 (217/413)	54.3 (176/324)
≥18 years	34.9 (144/413)	26.2 (85/324)
Missing	0.5 (2/413)	0.0 (0/324)
Lifetime sexual partners		
1	20.3 (84/413)	15.1 (49/324)
2	30.0 (124/413)	23.8 (77/324)
≥3	49.4 (204/413)	60.8 (197/324)
Missing	0.2 (1/413)	0.3 (1/324)
Used condoms during last sexual intercourse		
No	70.0 (289/413)	65.4 (212/324)
Yes	29.1 (120/413)	33.6 (109/324)
Missing	0.9 (4/413)	1.0 (3/324)
Method of contraceptive, last time had sex		
None/implant/ligation etc	51.1 (211/413)	52.2 (169/324)

(Continued)

Table 1. (Continued)

	Community clinic	Referral clinic
Variables	% (n/N)	% (n/N)
Injectables/birth control pill	21.1 (87/413)	20.7 (67/324)
Condoms	26.1 (108/413)	26.9 (87/324)
Missing	1.7 (7/413)	0.3 (1/324)
Using any contraception currently		
No	61.7 (255/413)	60.8 (197/324)
Yes	37.1 (153/413)	38.6 (125/324)
Missing	1.2 (5/413)	0.6 (2/324)
Pregnancy		
No	3.2 (13/413)	4.6 (15/324)
Yes	96.6 (399/413)	95.1 (308/324)
Missing	0.2 (1/413)	0.3 (1/324)

HIV: human immunodeficiency virus ARV's: Antiretrovirals, n: number of women responded, N: total number of study participants; Missing: not answered

<https://doi.org/10.1371/journal.pone.0241781.t001>

self-collected and clinician-collected samples was 91.4% [$k = 0.711$; 95%CI: 0.610 to 0.811, (Table 2)].

HR-HPV genotypes and agreement between clinician-collection and self-collection using the *hpVIR* assay

Of the 413 women attending the community-based clinic, 19.1% of the self-collected and 20.4% samples of clinician-collected samples were infected with single HR-HPV types, and this difference was not statistically significant ($p = 0.601$, Table 3). The prevalence of multiple HR-HPV infections was not significantly different between self-collected and clinician-collected specimens (8.7% vs 5.8%, $p = 0.108$). The most frequently detected HR-HPV genotypes were HPV16, HPV35, and the HPV33/52/58 group, both in self-collected and clinician-collected specimens (Fig 1). The agreement in the HR-HPV genotypes identified in self-collected and clinician-collected specimens ranged from moderate to almost perfect ($k = 0.571$ – 0.888), with HPV39 (99.8%, $k = 0.888$) and HPV31 (99.3%, $k = 0.838$) showing almost perfect agreement (Table 4).

Table 2. Concordance between high-risk HPV in self-collected and clinician-collected genital specimens from the community and the referral clinics.

Community clinic	Self-collected				
		HR-HPV positive (%)	HR-HPV negative (%)	Total (%)	% Agreement
Clinician-collected	HR-HPV positive (%)	85 (20.6)	24 (5.8)	109 (26.4)	86.9 ($k = 0.669$)
	HR-HPV negative (%)	30 (7.3)	274 (66.3)	304 (73.6)	
	Total (%)	115 (27.9)	298 (72.1)	413 (100.0)	
Referral clinic	Self-collected				
		HR-HPV positive (%)	HR-HPV negative (%)	Total (%)	% Agreement
Clinician-collected	HR-HPV positive (%)	251 (77.4)	20 (6.2)	271 (83.6)	91.4 ($k = 0.711$)
	HR-HPV negative (%)	8 (2.5)	45 (13.9)	53 (16.4)	
	Total (%)	259 (79.9)	65 (20.1)	324 (100.0)	

HR-HPV: high-risk human papillomavirus, k -kappa value

<https://doi.org/10.1371/journal.pone.0241781.t002>

Table 3. Comparison of self-collected and clinician-collected samples from women with single HR-HPV infections and multiple HR-HPV infections from the community and the referral clinics.

Variables	Community clinic			Referral clinic		
	Clinician-collected	Self-collected	p-value	Clinician-collected	Self-collected	p-value
Single infection	20.4% (85/413)	19.1% (79/413)	0.601	46.0% (149/324)	42.3% (137/324)	0.342
Multiple infection	5.8% (24/413)	8.7% (36/413)	0.108	37.7% (122/324)	37.7% (122/324)	1.000

<https://doi.org/10.1371/journal.pone.0241781.t003>

In the referral clinic, the distribution was similar when comparing HR-HPV genotypes in self-collected and clinician-collected samples. The HPV33/52/58 group, HPV16 and HPV35 were the most frequently detected HR-HPV genotypes in both collection groups (Fig 1). The prevalence of single HR-HPV infection was somewhat higher in clinician-collected specimens as compared to self-collected specimens, but the difference was not statistically significant [46.0% vs 42.3%, $p = 0.342$, (Table 3)]. The HR-HPV prevalence of multiple infection was the same in self-collected and clinician-collected specimens [37.7% vs 37.7%, $p = 1.000$, (Table 3)]. The concordance of the HR-HPV genotype distribution between collection groups showed kappa values ranging between 0.627–0.841 (Table 4). The HPV16 genotype showed a substantial agreement between self-collected and clinician-collected samples (89.2%, $k = 0.735$, Table 4), while HPV35 showed an almost perfect agreement between self-collected and clinician-collected specimens (95.4%, $k = 0.841$).

HR-HPV genotypes according to histology using *hpVIR* assay

Of the 324 women, histology results were available for 291 women including 2.1% (6/291) with inconclusive results (Table 5). More than half of the women had CIN3 (51.2%, 149/291), followed by CIN2 (23.0%, 67/291), no atypia (15.5%, 45/291), CIN1 (4.1%, 12/291) and CC (4.1%, 12/291). Overall, there was no significant difference in HR-HPV prevalence based on histology between self-collected and clinician-collected samples [78.4% (228/291) vs 81.8% (238/291), $p = 0.299$]. HR-HPV vaccine types (HPV16/18/31/33/45/52/58) were identified in 64.4% vs 46.7% of no atypia, 41.7% vs 41.7% of CIN1, 65.7% vs 59.7% of CIN2, 71.8% vs 71.1% of CIN3 and 75.0% vs 66.7% of CC between clinician-collected and self-collected, respectively (Table 5). The detection rate of the most commonly detected HPV genotypes (HPV16 and




Fig 1. Prevalence of HR-HPV genotypes in self-collected and clinician-collected specimens using the *hpVIR* assay in the community and the referral clinics.

<https://doi.org/10.1371/journal.pone.0241781.g001>

Table 4. Concordance for HR-HPV genotypes between self-collected and clinician-collected specimen in the community clinic and the referral clinics.

HR-HPV types	Community clinic			Referral clinic		
	% Agreement	kappa value	95% CI	% Agreement	Kappa value	95% CI
16	96.4	0.717	0.581–0.853	89.2	0.735	0.653–0.817
18/45	96.6	0.571	0.369–0.773	90.7	0.705	0.606–0.803
33/52/58	94.0	0.658	0.533–0.782	86.1	0.701	0.670–0.782
31	99.3	0.838	0.659–1.000	96.3	0.780	0.659–0.900
35	96.6	0.678	0.519–0.836	95.4	0.841	0.763–0.919
39	99.8	0.888	0.669–1.000	96.9	0.627	0.413–0.840
51	97.8	0.629	0.406–0.853	93.5	0.660	0.525–0.795
56	98.8	0.700	0.449–0.950	96.6	0.801	0.687–0.915
59	99.0	0.709	0.438–0.981	97.5	0.777	0.627–0.926

HR-HPV: high-risk human papillomavirus, CI: confidence intervals

<https://doi.org/10.1371/journal.pone.0241781.t004>

-35) was similar between self-collected and clinician-collected samples from women with CIN2/3 and CC [29.0% (66/228) vs 28.5% (65/228) and 19.7% (45/228) vs 19.3% (44/228), respectively]. Three women with CIN3 tested positive for HPV16 in their self-collected samples and were infected with multiple HPV types, but were negative in their clinician-collected samples. The HR-HPV copy number for three self-collected CIN3 samples were above the cut-off for HR-HPV positivity, but two of the samples had a low viral load.

Acceptance of self-collection of specimens for HPV testing

The participants from both study sites gave a similar response regarding self-collection versus clinician-collection of specimen (Table 6). More than half of the women find self-collection

Table 5. Distribution of HR-HPV types between self-collected and clinician-collected specimen according to histology results in women from referral clinic.

HR-HPV types	Histology results									
	NILM (n = 45)		CIN1 (n = 12)		CIN2 (n = 67)		CIN3 (n = 149)		Cervical cancer (n = 12)	
	Clinician-collected	Self-collected	Clinician-collected	Self-collected	Clinician-collected	Self-collected	Clinician-collected	Self-collected	Clinician-collected	Self-collected
16	11	10	0	2	18	17	41	44	6	5
18/45	7	8	1	1	16	19	23	29	3	3
33/52/58	12	9	2	3	19	19	66	62	3	2
31	5	3	2	2	4	6	18	15	0	0
35	6	6	2	1	14	15	29	28	1	2
39	2	0	1	0	2	2	5	8	0	0
51	1	4	1	1	4	7	17	16	1	1
56	4	3	1	1	5	7	12	14	2	2
59	4	3	0	0	4	4	6	13	1	1
HPV multiple infection	14	12	3	4	19	22	61	65	5	4
HR-HPV16/18/31/33/45/52/58	29	21	5	5	44	40	107	106	9	8
HR-HPV negative	13	17	5	5	14	15	20	25	1	1
HR-HPV positive	32	28	7	7	53	52	129	124	11	11

NILM: negative for intraepithelial lesions or malignancy, CIN: cervical intraepithelial neoplasia, HR-HPV: high-risk human papillomavirus

<https://doi.org/10.1371/journal.pone.0241781.t005>

Table 6. Experience of the mean at the with self-collection and clinician-collection.

Variable	Community clinic		Referral clinic	
	Clinician-collected %(n/N)	Self-collected %(n/N)	Clinician-collected %(n/N)	Self-collected %(n/N)
Embarrassed	12.8% (53/413)	14.0% (58/413)	16.4% (53/324)	18.2% (59/324)
Self-confident	17.0% (70/413)	16.0% (66/413)	16.7% (54/324)	16.7% (54/324)
Discomfort	3.9% (16/413)	3.4% (14/413)	9.5% (31/324)	4.3% (14/324)
Interested	64.4% (266/413)	66.1% (273/413)	54.0% (175/324)	59.3% (192/324)

<https://doi.org/10.1371/journal.pone.0241781.t006>

and clinician-collection to be interesting (64.4% vs 66.1% in the community-based clinic and 54.0% vs 59.3% in referral clinic, respectively). A very small proportion of women from the community-based clinic experienced discomfort when collecting the sample, both using self-collection (3.4%) and clinician-collection (3.9%). In the referral clinic, a slightly higher proportion of women experienced discomfort with clinician-collection (9.5%) as compared to self-collection (4.3%) of samples. At both study sites older women reported to be embarrassed and were more likely to have had only one Pap smear screening test in their lifetime [73.3% (22/30) self-collected vs 66.7% (20/30) clinician-collected at community-based clinic and 66.1% (39/59) self-collected vs 67.9% (36/53) clinician-collected at referral clinic]. Most participants at both study sites reported that they would be willing to perform self-collection at home and return the card for testing [77.2% in the community-based clinic and 83.0% in referral clinic, (Table 7)]. Given a choice, almost all participants preferred clinician-collection to self-collection of samples (95.6% in community-based clinic and 90.7% in referral clinic, Table 7).

HPV detection in cervical clinician-collected specimen using the *hpVIR* assay and HC-2

A total of 628 women were screened for HR-HPV infection using HC-2 and the *hpVIR* assay. Overall, the HR-HPV prevalence was 46.2% using *hpVIR* compared to 48.3% using HC-2, with an agreement of 87.7% ($k = 0.754$, 95%CI:0.703–0.806) between the two assays (Table 8).

Table 7. Investigation of the acceptability of self-collection for HPV testing.

	Community clinic	Referral clinic
Variables	% (n/N)	% (n/N)
Would you be willing to collect a sample yourself (self-sample) at home and bring it in for testing?		
Yes	77.2% (319/413)	83.0% (269/324)
Not sure	18.6% (77/413)	10.8% (35/324)
No	1.9% (8/413)	4.3% (14/324)
If no, why not?		
I prefer the specimen to be taken by the nurse	12.5% (1/8)	12.5% (1/14)
I will not be able to take the specimen correctly	62.5% (5/8)	62.5% (5/14)
Which method do you prefer for cervical cancer screening		
To take a sample myself	1.4% (6/413)	3.7% (12/324)
For a healthcare worker to take a sample	95.6% (395/413)	90.7% (294/324)
Either for myself to take a sample or a health worker to take a sample	1.2% (5/413)	2.5% (8/324)

<https://doi.org/10.1371/journal.pone.0241781.t007>

Table 8. Concordance of HR-HPV infection between the *hpVIR* and the HC-2 assays.

HC-2		<i>hpVIR</i>			% Agreement
		HR-HPV positive (%)	HR-HPV negative (%)	Total (%)	
	HR-HPV positive (%)	258 (41.1)	45 (7.1)	303 (48.3)	87.7 (k = 0.754)
	HR-HPV negative (%)	32 (5.1)	293 (46.7)	325 (51.7)	
	Total (%)	290 (46.2)	338 (53.8)	628 (100.0)	

HR-HPV: high-risk human papillomavirus, k-kappa value

<https://doi.org/10.1371/journal.pone.0241781.t008>

Discussion

The present study assessed the prevalence of HR-HPV genotypes between self-collected and clinician-collected samples applied to FTA cards in women from a community-based clinic and a referral clinic in Eastern Cape, South Africa. We observed high HR-HPV prevalence, overall agreement of HR-HPV infection, and a similar distribution of HR-HPV genotypes between self-collected and clinician-collected samples. A similar trend has been reported using FTA cards and PCR-based assays performed in women from the general population and referral population [23, 26, 31]. Our findings confirm that the self-collection method and storage on FTA cards is an adequate procedure for HPV DNA testing and support the use of self-collection as an alternative strategy in the regular CC screening routine. Many developed countries have transitioned to HPV DNA testing, and CC screening guidelines with HPV DNA testing has been implemented [17, 18, 32]. However, this is the opposite for low-income countries having low participation rate in CC screening particularly in rural regions. Therefore, introducing new screening strategies for women experiencing screening barriers will improve uptake and could have an impact on the incidence of CC diseases [33].

The prevalence of HR-HPV infection and distribution of HR-HPV genotypes differs between regions and countries [1, 34]. In the current study, HPV16 and HPV35 were the most frequently detected genotypes in both self-collected and clinician-collected samples in the two sites. Other cross-sectional studies conducted in South Africa also reported HPV35 and HPV16 as the most commonly identified genotypes among women with or without cervical abnormalities [35–37]. However, the predominate HR-HPV genotypes in other African studies were HPV18 & HPV59 in Ghana and HPV52 & HPV16 in Tanzania [35, 38]. A recent study done in Brazil among women residing in rural areas showed HPV56 and HPV51 the common types in their population [39]. The two HR-HPV genotypes (HPV16 and 35) are among the top five causing CC in South Africa [3]. None of the HPV vaccines protect against HPV35 (overall prevalence = 18.5%, mostly in multiple infections); an important HR-HPV genotype in women with cervical disease in the Eastern Cape. A high proportion of 72.0% CIN3/CC women were more likely to harbour vaccine HR-HPV types (HPV16/18/31/33/45/52/58). Our results indicate that studies on the distribution of HPV genotypes in CC not present in the current vaccines should be considered when developing strategies and targets for the next generation of HPV vaccines.

A slightly higher prevalence of multiple HR-HPV infections was observed in self-collected as compared to clinician-collected samples in the community-based clinic which could reflect the anatomic area sampled. Vaginal self-collected samples contain vaginal fluid with both cervical and vaginal cells. Women are more likely to be infected with several HPV genotypes in the vagina that are not present in the endocervix [22, 31]. HR-HPV genotypes at low viral load present only in vaginal self-collected samples are not correlated with CC diseases [22, 40]. The few cases of discordance of HPV16 between self-collected and clinician-collected samples among women with CIN3 in our study, were likely to be due to low viral load.

There was high concordance in HR-HPV detection between self-collected and clinician-collected samples in the present study. The use of the Viba-brush in combination with FTA cards, yields good quality DNA in both self-collected and clinician-collected specimen, indicating that the self-collection was performed correctly. Sampling devices for HPV DNA testing have a strong effect on the detection of HPV infection. Self-sampling with a brush-based sampling device in combination with FTA cards was found to be acceptable among women and this together with the clinically validated PCR-based HPV DNA test yielded high agreement in identifying women with high risk of CIN2+ in comparison with clinician-collection sampling [31]. The in-house *hpVIR* assay combined with the FTA cards performed well compared to HC-2, as demonstrated by substantial agreement in HR-HPV positivity. Our findings concur with previous studies reporting on the performance of the two assays, indicating that the *hpVIR* assay and FTA cards can be utilised for primary screening programmes in regions other than Sweden [29, 30].

At both study sites, women showed the same attitude regarding self-collection versus clinician collection samples for HPV DNA testing. The high rate of willingness to perform self-collection was not influenced by sociodemographic factors or any other factor studied as observed in previous studies [41, 42]. This underscores the strong scientific basis for implementing self-collection in CC screening, thereby enabling women from communities and rural settings where CC screening programmes are presently not available to participate. Moreover, the high rate of acceptability for vaginal self-collection could influence the effectiveness of HPV self-collection in routine screening programme [43]. Although self-collection was highly accepted, the majority of women preferred the specimen to be taken by a healthcare worker. In our study, we did not investigate factors influencing women's choice as to why they prefer clinician-collection of samples. Mao and colleagues (2017) reported that women who prefer a clinician-collection want to have a one-on-one consultation with a clinician, so that they can address other issues they may have [44]. Other factors that women are raising include adequacy of the specimen and lack of confidence to perform self-collection correctly [42, 45]. Therefore, our findings suggest that women should be given an opportunity to choose which method they prefer for specimen collection, when participating in the CC screening programme.

A large population of women in the Eastern Cape Province reside in rural areas and are less likely to participate in CC screening programme because of the lack of nearby facilities with functional CC screening programmes [7]. Implementing self-sampling in this region will therefore have an impact on the screening programme since it does not require infrastructure. This approach allows women who are unwilling to attend Pap smear test performed by clinician to have access to cervical screening and give them privacy to collect the specimen [23]. Implementation of self-collection is also more likely to create awareness and more information about HPV infection than clinician-collection [46, 47]. There are different methods to administer the self-collection kit, including mailing the kit directly to the woman, conducting community-based campaigns and using an opt-in strategy [19]. In developed countries, such as Sweden, mailing self-collection kits to underscreened women has been shown to work effectively [33]. However, in South Africa, it will be challenging to send self-collection kits as the postal system is not working effectively or does not exist in some rural areas. Direct offer of self-collection kit would be a better method to achieve knowledge and confidence among women residing in rural communities as this method has been previously shown to have significantly high rate (>75%) among underscreened women [19]. A second issue would be loss of women to follow-up due to difficulty in travelling to the clinics, something which would have a strong negative impact on the efficiency of CC screening programmes. Therefore, effective communication strategies to follow-up women with HR-HPV positive test needs to be

developed prior to implementation of self-collection for HPV testing outside the clinics in South Africa.

The FTA cards provide several advantages for self-collection in this environment. These strategies have the potential to improve CC screening programmes by increasing the screening coverage, reducing the work load for the clinicians, and identify women at high-risk of developing CC [19, 33]. Furthermore, this allows healthcare facilities with limited resources for CC screening to target women at high-risk of cervical cancer. However, self-collection for HPV has been found to increase the rate of women being referred to colposcopy clinics and for treatment. Biomarkers such as DNA methylation has been reported to triage self-collected HR-HPV positive women, but further studies are needed to determine its clinical value [48]. Repeating the HPV test in 4–6 months for women that are HPV positive in their screening test, can be used to identify women with persistent infection and thereby increase the sensitivity and specificity of screening [33]. If it was possible to rescreen women this may be the best approach to triage women for referral to the treatment clinic.

A potential limitation in this study could be the use of in-house *hpVIR* real time PCR assay. The *hpVIR* assay is clinically validated HPV test that gives HPV type information, but it does not distinguish between the alpha HPV types, HPV18/45 and HPV33/52/58 because they are closely related. This limits the ability to provide the prevalence of the individual HPV types that are grouped in the *hpVIR* assay.

Conclusion

The study shows a high concordance in HR-HPV genotype prevalence between self-collected and clinician-collected specimens. The frequency distribution of HR-HPV genotypes will assist in identifying and monitoring genotypes that are not covered by the vaccines currently in use. The self-collection procedure for HPV testing was regarded as highly acceptable by the women. Self-collection can have a positive impact on the cervical screening programme in South Africa by increasing the population coverage in rural areas, and enable women who are unable to attend clinics to participate in cervical cancer screening.

Supporting information

S1 Data.

(CSV)

S2 Data.

(CSV)

Acknowledgments

We thank the Community clinic staff members, Sr Virginia Maqoga, Luviwe Lutotswana and women who kindly participated in the study.

Meeting information

Part of the data was presented at the Virology Africa Conference, Cape Town, 10–14 February 2020.

Author Contributions

Conceptualization: Tracy L. Meiring, Zizipho Z. A. Mbulawa, Charles B. Businge, Ulf Gyllensten, Anna-Lise Williamson.

Data curation: Ongeziwe Taku, Inger Gustavsson, Keletso Phohlo, Mirta Garcia-Jardon, Zizipho Z. A. Mbulawa.

Formal analysis: Ongeziwe Taku, Inger Gustavsson, Mirta Garcia-Jardon.

Funding acquisition: Zizipho Z. A. Mbulawa, Charles B. Businge, Ulf Gyllensten, Anna-Lise Williamson.

Investigation: Ongeziwe Taku, Tracy L. Meiring, Keletso Phohlo, Mirta Garcia-Jardon, Zizipho Z. A. Mbulawa, Charles B. Businge, Ulf Gyllensten, Anna-Lise Williamson.

Methodology: Ongeziwe Taku, Inger Gustavsson, Zizipho Z. A. Mbulawa, Charles B. Businge, Ulf Gyllensten, Anna-Lise Williamson.

Project administration: Ongeziwe Taku, Inger Gustavsson, Zizipho Z. A. Mbulawa, Charles B. Businge, Ulf Gyllensten, Anna-Lise Williamson.

Resources: Zizipho Z. A. Mbulawa, Charles B. Businge, Ulf Gyllensten, Anna-Lise Williamson.

Software: Inger Gustavsson.

Supervision: Tracy L. Meiring, Inger Gustavsson, Zizipho Z. A. Mbulawa, Charles B. Businge, Ulf Gyllensten, Anna-Lise Williamson.

Validation: Inger Gustavsson.

Visualization: Ongeziwe Taku.

Writing – original draft: Ongeziwe Taku.

Writing – review & editing: Ongeziwe Taku, Tracy L. Meiring, Inger Gustavsson, Keletso Phohlo, Mirta Garcia-Jardon, Zizipho Z. A. Mbulawa, Charles B. Businge, Ulf Gyllensten, Anna-Lise Williamson.

References

1. Bruni L, Albero G, Serrano B, Mena M, Gómez D, Muñoz J, et al. 2019. CO/IARC Information Centre on HPV and Cancer (HPV Information Centre). Human Papillomavirus and Related Diseases in the World. Summary Report <https://www.hpvcentre.net/statistics/reports/XWX.pdf> (Accessed 28 June 2019).
2. Arbyn M, Weiderpass E, Bruni L, de Sanjosé S, Saraiya M, Ferlay J, et al. 2020. Estimates of incidence and mortality of cervical cancer in 2018: a worldwide analysis. *Lancet Glob Health* 8:e191–e203. [https://doi.org/10.1016/S2214-109X\(19\)30482-6](https://doi.org/10.1016/S2214-109X(19)30482-6) PMID: 31812369
3. Bruni L, Albero G, Serrano B, Mena M, Gómez D, Muñoz J, et al. 2019. ICO/IARC Information Centre on HPV and Cancer (HPV Information Centre). Human Papillomavirus and Related Diseases in South Africa. Summary Report <https://hpvcentre.net/statistics/reports/ZAF.pdf> (Accessed 28 February 2019).
4. Africa NfcdoS. 2020. Summary statistics of cancer diagnosed histologically in 2016. https://www.nicd.ac.za/wp-content/uploads/2020/04/NCR_2016_Report_updated_14April2020.pdf (Accessed June 2020).
5. Arbyn M, Xu L, Simoens C, Martin-Hirsch PP. 2018. Prophylactic vaccination against human papillomaviruses to prevent cervical cancer and its precursors. *Cochrane Database Syst Rev* 5: Cd009069. <https://doi.org/10.1002/14651858.CD009069.pub3> PMID: 29740819
6. Denny L, Herrero R, Levin C, Kim JJ. 2015. Cervical Cancer. In Gelband H, Jha P, Sankaranarayanan R, Horton S (ed), *Cancer: Disease Control Priorities, Third Edition (Volume 3)* 10.1596/978-1-4648-0349-9_ch4. The International Bank for Reconstruction and Development / The World Bank © 2015 International Bank for Reconstruction and Development / The World Bank., Washington (DC).
7. Hoque M, Hoque E, Kader SB. 2008. Evaluation of cervical cancer screening program at a rural community of South Africa. *East Afr J Public Health* 5:111–6. PMID: 19024420
8. WHO. 2014. Comprehensive cervical cancer control—A guide to essential practice. <https://www.who.int/reproductivehealth/publications/cancers/cervical-cancer-guide/en/> (accessed March 2020).

9. Makura CBT, Schnippel K, Michelow P, Chibwasha CJ, Goeieman B, Jordaan S, et al. 2016. Choropleth Mapping of Cervical Cancer Screening in South Africa Using Healthcare Facility-level Data from the National Laboratory Network. *AIMS public health* 3:849–862. <https://doi.org/10.3934/publichealth.2016.4.849> PMID: 29546199
10. Sibiya MN, Grainger L. 2007. An assessment of the implementation of the provincial cervical screening programme in selected Primary Health Care Clinics in the Ilembe Region, KwaZulu-Natal. *Curatationis* 30:48–55. <https://doi.org/10.4102/curatationis.v30i1.1050> PMID: 17515316
11. Nene B, Jayant K, Arrossi S, Shastri S, Budukh A, Hingmire S, et al. 2007. Determinants of womens participation in cervical cancer screening trial, Maharashtra, India. *Bull World Health Organ* 85:264–72. <https://doi.org/10.2471/blt.06.031195> PMID: 17546307
12. Waller J, Bartoszek M, Marlow L, Wardle J. 2009. Barriers to cervical cancer screening attendance in England: a population-based survey. *J Med Screen* 16:199–204. <https://doi.org/10.1258/jms.2009.009073> PMID: 20054095
13. Daley E, Alio A, Anstey EH, Chandler R, Dyer K, Helmy H. 2011. Examining barriers to cervical cancer screening and treatment in Florida through a socio-ecological lens. *J Community Health* 36:121–31. <https://doi.org/10.1007/s10900-010-9289-7> PMID: 20559695
14. Ogilvie GS, van Niekerk D, Kraijden M, Smith LW, Cook D, Gondara L, et al. 2018. Effect of Screening With Primary Cervical HPV Testing vs Cytology Testing on High-grade Cervical Intraepithelial Neoplasia at 48 Months: The HPV FOCAL Randomized Clinical Trial. *Jama* 320:43–52. <https://doi.org/10.1001/jama.2018.7464> PMID: 29971397
15. Kitchener H. 2019. Optimising future cervical screening strategies. *Papillomavirus Res* 7:201–203. <https://doi.org/10.1016/j.pvr.2019.04.001> PMID: 30978414
16. Sankaranarayanan R, Nene BM, Shastri SS, Jayant K, Muwonge R, Budukh AM, et al. 2009. HPV screening for cervical cancer in rural India. *N Engl J Med* 360:1385–94. <https://doi.org/10.1056/NEJMoa0808516> PMID: 19339719
17. Polman NJ, de Haan Y, Veldhuijzen NJ, Heideman DAM, de Vet HCW, Meijer C, et al. 2019. Experience with HPV self-sampling and clinician-based sampling in women attending routine cervical screening in the Netherlands. *Prev Med* 125:5–11. <https://doi.org/10.1016/j.ypmed.2019.04.025> PMID: 31054907
18. Committee MSA. 2014. National Cervical Screening Program renewal: effectiveness modelling and economic evaluation in the Australian setting. Report November 2013. MSAC application 1276 Canberra: Australian Government Department of Health [http://www.cancerscreening.gov.au/internet/screening/publishingnsf/Content/E6A211A6FFC29E2CCA257CED007FB678/\\$File/Renewal%20Economic%20Evaluation.pdf](http://www.cancerscreening.gov.au/internet/screening/publishingnsf/Content/E6A211A6FFC29E2CCA257CED007FB678/$File/Renewal%20Economic%20Evaluation.pdf).
19. Arbyn M, Smith SB, Temin S, Sultana F, Castle P. 2018. Detecting cervical precancer and reaching underscreened women by using HPV testing on self samples: updated meta-analyses. *Bmj* 363:k4823.
20. Gök M, Heideman DA, van Kemenade FJ, Berkhof J, Rozendaal L, Spruyt JW, et al. 2010. HPV testing on self collected cervicovaginal lavage specimens as screening method for women who do not attend cervical screening: cohort study. *Bmj* 340:c1040. <https://doi.org/10.1136/bmj.c1040> PMID: 20223872
21. Tranberg M, Bech BH, Blaakær J, Jensen JS, Svanholm H, Andersen B. 2018. Preventing cervical cancer using HPV self-sampling: direct mailing of test-kits increases screening participation more than timely opt-in procedures—a randomized controlled trial. *BMC Cancer* 18:273. <https://doi.org/10.1186/s12885-018-4165-4> PMID: 29523108
22. Gustavsson I, Sanner K, Lindell M, Strand A, Olovsson M, Wikström I, et al. 2011. Type-specific detection of high-risk human papillomavirus (HPV) in self-sampled cervicovaginal cells applied to FTA elute cartridge. *J Clin Virol* 51:255–8. <https://doi.org/10.1016/j.jcv.2011.05.006> PMID: 21632283
23. Guan Y, Gravitt PE, Howard R, Eby YJ, Wang S, Li B, et al. 2013. Agreement for HPV genotyping detection between self-collected specimens on a FTA cartridge and clinician-collected specimens. *J Virol Methods* 189:167–71. <https://doi.org/10.1016/j.jviromet.2012.11.010> PMID: 23370404
24. Gustavsson I, Lindell M, Wilander E, Strand A, Gyllensten U. 2009. Use of FTA card for dry collection, transportation and storage of cervical cell specimen to detect high-risk HPV. *J Clin Virol* 46:112–6. <https://doi.org/10.1016/j.jcv.2009.06.021> PMID: 19628427
25. de Bie RP, Schmeink CE, Bakkers JM, Snijders PJ, Quint WG, Massuger LF, et al. 2011. The indicating FTA elute cartridge a solid sample carrier to detect high-risk HPV and high-grade cervical lesions. *J Mol Diagn* 13:371–6. <https://doi.org/10.1016/j.jmoldx.2011.02.003> PMID: 21704269
26. Geraets DT, van Baars R, Alonso I, Ordi J, Torné A, Melchers WJ, et al. 2013. Clinical evaluation of high-risk HPV detection on self-samples using the indicating FTA-elute solid-carrier cartridge. *J Clin Virol* 57:125–9. <https://doi.org/10.1016/j.jcv.2013.02.016> PMID: 23518442
27. Lenselink CH, de Bie RP, van Hamont D, Bakkers JM, Quint WG, Massuger LF, et al. 2009. Detection and genotyping of human papillomavirus in self-obtained cervicovaginal samples by using the FTA

- cartridge: new possibilities for cervical cancer screening. *J Clin Microbiol* 47:2564–70. <https://doi.org/10.1128/JCM.00285-09> PMID: 19553570
28. Sellors J, editors. 2003. Colposcopy and treatment if cervical intraepithelial neoplasia: a beginners' manual. IARC; <https://screeningiarcrf/colpophp>.
 29. Gustavsson I, Aarnio R, Myrnäs M, Hedlund-Lindberg J, Taku O, Meiring T, et al. 2019. Clinical validation of the HPVIR high-risk HPV test on cervical samples according to the international guidelines for human papillomavirus DNA test requirements for cervical cancer screening. *Virology* 16:107. <https://doi.org/10.1186/s12985-019-1216-7> PMID: 31438976
 30. Gustavsson I, Juko-Pecirep I, Backlund I, Wilander E, Gyllensten U. 2009. Comparison between the Hybrid Capture 2 and the hpVIR real-time PCR for detection of human papillomavirus in women with ASCUS or low grade dysplasia. *J Clin Virol* 45:85–9. <https://doi.org/10.1016/j.jcv.2009.04.012> PMID: 19451022
 31. Dijkstra MG, Heideman DA, van Kemenade FJ, Hogewoning KJ, Hesselink AT, Verkuijten MC, et al. 2012. Brush-based self-sampling in combination with GP5+/6+-PCR-based hrHPV testing: high concordance with physician-taken cervical scrapes for HPV genotyping and detection of high-grade CIN. *J Clin Virol* 54:147–51. <https://doi.org/10.1016/j.jcv.2012.02.022> PMID: 22445557
 32. Arbyn M, Haelens A, Desomer A, Verdoordt F, Thiry N, Francart J, et al. 2015. Cervical cancer screening program and Human Papillomavirus (HPV) testing, part II: Update on HPV primary screening. Health Technology Assessment (HTA) Brussels: Belgian Health Care Knowledge Centre (KCE) KCE Reports 238.
 33. Gustavsson I, Aarnio R, Berggrund M, Hedlund-Lindberg J, Strand AS, Sanner K, et al. 2018. Randomised study shows that repeated self-sampling and HPV test has more than two-fold higher detection rate of women with CIN2+ histology than Pap smear cytology. *Br J Cancer* 118:896–904. <https://doi.org/10.1038/bjc.2017.485> PMID: 29438367
 34. de Sanjosé S, Diaz M, Castellsagué X, Clifford G, Bruni L, Muñoz N, Bosch FX. 2007. Worldwide prevalence and genotype distribution of cervical human papillomavirus DNA in women with normal cytology: a meta-analysis. *Lancet Infect Dis* 7:453–9. [https://doi.org/10.1016/S1473-3099\(07\)70158-5](https://doi.org/10.1016/S1473-3099(07)70158-5) PMID: 17597569
 35. Dols JAM, Reid G, Brown JM, Tempelman H, Bontekoe TR, Quint WGV, et al. 2012. HPV Type Distribution and Cervical Cytology among HIV-Positive Tanzanian and South African Women. *ISRN Obstetrics and Gynecology* 2012:514146. <https://doi.org/10.5402/2012/514146> PMID: 22811925
 36. McDonald AC, Denny L, Wang C, Tsai W-Y, Wright TC Jr., Kuhn L. 2012. Distribution of high-risk human papillomavirus genotypes among HIV-negative women with and without cervical intraepithelial neoplasia in South Africa. *PloS one* 7:e44332–e44332. <https://doi.org/10.1371/journal.pone.0044332> PMID: 22970201
 37. McDonald A, Tergas A, Kuhn L, Denny L, Wright T. 2014. Distribution of Human Papillomavirus Genotypes among HIV-Positive and HIV-Negative Women in Cape Town, South Africa. *Frontiers in oncology* 4:48. <https://doi.org/10.3389/fonc.2014.00048> PMID: 24672770
 38. Awua AK, Sackey ST, Osei YD, Asmah RH, Wiredu EK. 2016. Prevalence of human papillomavirus genotypes among women with cervical cancer in Ghana. *Infectious agents and cancer*. 11(4). <https://doi.org/10.1186/s13027-016-0050-4> PMID: 26816527
 39. Lorenzi AT, Fregnani JH, Villa LL, Sichero L, Nunes EM, Longatto-Filho A. 2019. Diversity of human papillomavirus typing among women population living in rural and remote areas of Brazilian territory. *Papillomavirus Res* 8:100186. <https://doi.org/10.1016/j.pvr.2019.100186> PMID: 31539646
 40. Zhang SK, Ci PW, Velicer C, Kang LN, Liu B, Cui JF, et al. 2014. Comparison of HPV genotypes and viral load between different sites of genital tract: the significance for cervical cancer screening. *Cancer Epidemiol* 38:168–73. <https://doi.org/10.1016/j.canep.2014.01.004> PMID: 24522235
 41. Murchland AR, Gottschlich A, Bevilacqua K, Pineda A, Sandoval-Ramírez BA, Alvarez CS, et al. 2019. HPV self-sampling acceptability in rural and indigenous communities in Guatemala: a cross-sectional study. *BMJ Open* 9:e029158. <https://doi.org/10.1136/bmjopen-2019-029158> PMID: 31662358
 42. Morgan K, Azzani M, Khaing SL, Wong YL, Su TT. 2019. Acceptability of Women Self-Sampling versus Clinician-Collected Samples for HPV DNA Testing: A Systematic Review. *J Low Genit Tract Dis* 23:193–199. <https://doi.org/10.1097/LGT.0000000000000476> PMID: 30933030
 43. Polman NJ, Ebisch RMF, Heideman DAM, Melchers WJG, Bekkers RLM, Molijn AC, et al. 2019. Performance of human papillomavirus testing on self-collected versus clinician-collected samples for the detection of cervical intraepithelial neoplasia of grade 2 or worse: a randomised, paired screen-positive, non-inferiority trial. *Lancet Oncol* 20:229–238.
 44. Mao C, Kulasingam SL, Whitham HK, Hawes SE, Lin J, Kiviati NB. 2017. Clinician and Patient Acceptability of Self-Collected Human Papillomavirus Testing for Cervical Cancer Screening. *J Womens Health (Larchmt)* 26:609–615. <https://doi.org/10.1089/jwh.2016.5965> PMID: 28332888

45. Saidu R, Moodley J, Tergas A, Momberg M, Boa R, Wright T, et al. 2018. South African women's perspectives on self-sampling for cervical cancer screening: A mixed-methods study. *S Afr Med J* 109:47–52. <https://doi.org/10.7196/SAMJ.2018.v109i1.13278> PMID: 30606304
46. Allen-Leigh B, Uribe-Zúñiga P, León-Maldonado L, Brown BJ, Lőrincz A, Salmeron J, et al. 2017. Barriers to HPV self-sampling and cytology among low-income indigenous women in rural areas of a middle-income setting: a qualitative study. *BMC Cancer* 17:734. <https://doi.org/10.1186/s12885-017-3723-5> PMID: 29121873
47. Crofts V, Flahault E, Tebeu PM, Untiet S, Fosso GK, Boulvain M, et al. 2015. Education efforts may contribute to wider acceptance of human papillomavirus self-sampling. *Int J Womens Health* 7:149–54. <https://doi.org/10.2147/IJWH.S56307> PMID: 25674016
48. Snellenberg S, De Strooper LM, Hesselink AT, Meijer CJ, Snijders PJ, Heideman DA, et al. 2012. Development of a multiplex methylation-specific PCR as candidate triage test for women with an HPV-positive cervical scrape. *BMC Cancer* 12:551. <https://doi.org/10.1186/1471-2407-12-551> PMID: 23176198