

Comparison of amplicor and GeneXpert MTB/ RIF tests for diagnosis of tuberculous meningitis.

| | |
|---------------|---|
| Item Type | Article |
| Authors | Patel, Vinod B;Connolly, Cathy;Singh, Ravesh;Lenders, Laura;Matinyenya, Brian;Theron, Grant;Ndung'u, Thumbi;Dheda, Keertan |
| Citation | Comparison of amplicor and GeneXpert MTB/RIF tests for diagnosis of tuberculous meningitis. 2014, 52 (10):3777-80 J. Clin. Microbiol. |
| DOI | 10.1128/JCM.01235-14 |
| Journal | Journal of clinical microbiology |
| Rights | Archived with thanks to Journal of clinical microbiology |
| Download date | 2024-08-06 23:31:58 |
| Link to Item | http://hdl.handle.net/11288/595134 |

Comparison of Amplicor and GeneXpert MTB/RIF Tests for Diagnosis of Tuberculous Meningitis

Vinod B. Patel, Cathy Connolly, Ravesh Singh, Laura
Lenders, Brian Matinyenya, Grant Theron, Thumbi Ndung'u
and Keertan Dheda

J. Clin. Microbiol. 2014, 52(10):3777. DOI:
10.1128/JCM.01235-14.

Published Ahead of Print 23 July 2014.

Updated information and services can be found at:
<http://jcm.asm.org/content/52/10/3777>

SUPPLEMENTAL MATERIAL

These include:

[Supplemental material](#)

REFERENCES

This article cites 28 articles, 12 of which can be accessed free
at: <http://jcm.asm.org/content/52/10/3777#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new
articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Comparison of Amplicor and GeneXpert MTB/RIF Tests for Diagnosis of Tuberculous Meningitis

Vinod B. Patel,^a Cathy Connolly,^b Ravesh Singh,^{c,d} Laura Lenders,^e Brian Matinyenya,^e Grant Theron,^e Thumbi Ndung'u,^{c,d} Keertan Dheda^{e,f,*}

Department of Neurology, University of KwaZulu-Natal, Durban, South Africa^a; Biostatistics Unit, Medical Research Council, Durban, South Africa^b; KwaZulu-Natal Research Institute for Tuberculosis and HIV (K-RITH), Nelson R. Mandela School of Medicine, University of KwaZulu-Natal, Durban, South Africa^c; HIV Pathogenesis Programme, Doris Duke Medical Research Institute, Nelson R. Mandela School of Medicine, University of KwaZulu-Natal, Durban, South Africa^d; Lung Infection and Immunity Unit, Division of Pulmonology and UCT Lung Institute, Department of Medicine, University of Cape Town, Cape Town, South Africa^e; Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town, South Africa^f

There are no data about the comparative accuracy of commercially available nucleic acid amplification tests (GeneXpert MTB/RIF and Roche Amplicor) for the diagnosis of tuberculous meningitis (TBM). A total of 148 patients with suspected TBM were evaluated, and cultures served as the reference standard. The sensitivities and specificities (95% confidence interval [CI]) for the Amplicor and Xpert MTB/RIF tests were similar: 46 (31–60) versus 50 (33–67) and 99 (93–100) and 94 (84–99), respectively.

There are ~10 million new cases and 1.7 million deaths from tuberculosis (TB) annually (1). In sub-Saharan Africa and particularly South Africa, up to 80% of TB cases are HIV coinfecting (1) and approximately 40% have extrapulmonary tuberculosis (EPTB). One-tenth of these have tuberculous meningitis (TBM) (2, 3). Patients with TBM frequently require prolonged admission to hospitals and have high morbidity rates due to neuropathology with substantial mortality (~30%), particularly if the diagnosis and follow-on therapy are delayed (4–7). Thus, rapid diagnosis of TBM is essential for early institution of the appropriate therapy. However, the current tools such as smear microscopy perform very poorly in TBM. A systematic review published more than a decade ago showed that nucleic acid amplification tests (NAATs) had an overall sensitivity of 56% and specificity approaching 100% (8).

More recently, however, more sensitive platforms have become available. The Xpert MTB/RIF is a new cartridge-based real-time heminested closed NAAT platform, presently being rolled out in resource-poor settings as a potential easy-to-use point-of-care test (9). The Amplicor PCR is an alternative closed NAAT platform that is now also commercially available (10). We recently reported our experience with Xpert MTB/RIF in TBM (11). However, there are no comparative data evaluating its accuracy in TBM. Here we report our experience using the Xpert MTB/RIF and Amplicor assays side by side, employing cultures as the reference standard.

Methods are outlined only briefly here. More detailed methods are given in the supplement material. A total of 148 consecutive patients with suspected meningitis were prospectively recruited between January 2008 and December 2010. Patients with a meningitic illness who were referred from local district general hospitals were investigated at Inkosi Albert Luthuli Central Hospital (IALCH), a tertiary hospital. Patients had a computed tomography (CT) scan done to exclude contraindications to a lumbar puncture, and blood samples were collected for routine tests, including HIV infection and a CD4 count, and for exclusion of alternate causes of meningitis, including serum fluorescent treponemal antibody (FTA) and Venereal Disease Research Laboratory (VDRL) tests, as previously outlined (12). Cerebrospinal fluid (CSF) was processed for the following tests: microscopy (Gram stain and fluorescence staining for acid-fast bacilli [auramine]), bacterial culture, *Mycobacterium tuberculosis* culture (Bactec 960

MGIT; Becton Dickinson Diagnostic Systems, Sparks, MD), fungal culture, and the cryptococcal latex agglutination test (CLAT). In addition to routine testing, CSF was processed for the Amplicor PCR *Mycobacterium tuberculosis* test (Roche Diagnostic Systems Inc., Branchburg, NJ) (Amplicor PCR) and the Xpert MTB/RIF test (Cepheid, Sunnyvale CA, USA).

Recently archived (–70°C) and uncentrifuged samples ($n = 148$) were processed in an independent laboratory using the Amplicor kit for the detection of *M. tuberculosis*. These samples were used in a previous publication evaluating the Xpert MTB/RIF test for the diagnosis of TBM (11). Here we report on the head-to-head comparison of the two PCR assays, which had not previously been reported. The Amplicor test was done as per the manufacturer's protocol. Briefly, 0.5 ml of CSF was used to extract DNA with the Roche MagNA Pure automated DNA extraction system using a high-performance DNA isolation kit. The extracted DNA was then amplified using the biotinylated primers KY18 and KY75 as described in the kit protocol (Roche Amplicor *Mycobacterium tuberculosis* test). PCR products were detected by the Cobas Amplicor analyzer according to the kit protocol. The samples were also processed for Xpert MTB/RIF analysis at the Lung Infection and Immunity Unit Laboratory (Department of Medicine, Groote Schuur Hospital, University of Cape Town), according to the manufacturer's instructions (13). The laboratory technicians performing the Amplicor PCR and Xpert MTB/RIF assays were blinded to all subject details.

Received 30 April 2014 Returned for modification 5 June 2014

Accepted 14 July 2014

Published ahead of print 23 July 2014

Editor: G. A. Land

Address correspondence to Keertan Dheda, keertan.dheda@uct.ac.za.

* Present address: Keertan Dheda, Department of Medicine, Groote Schuur Hospital Observatory, Cape Town, South Africa.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/JCM.01235-14>.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JCM.01235-14




FIG 1 Summary of the study plan, sample processing, and outcome. +ve, positive; -ve, negative.

Patients were categorized, based on standardized published diagnostic criteria, as definite TBM if the CSF *M. tuberculosis* culture and/or the Amplicor PCR test was positive (14, 15), probable TBM (treated empirically with anti-TB drugs but not meeting the definite TBM criteria), or non-TBM (alternate diagnosis confirmed and response to therapy documented in the absence of anti-TB treatment) (16).

The characteristics of definite TBM and non-TBM patients were compared using the chi square test or Fisher's exact test for categorical variables and Wilcoxon's rank sum test for continuous variables. Sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV), overall agreement, and likelihood ratios are reported as measures of diagnostic efficacy. Specificity and sensitivity between the Amplicor PCR and Xpert MTB/RIF tests were compared using McNemar's chi square test. Data were analyzed using Stata v12 (Statacorp, USA).

Figure 1 outlines the study plan and summarizes the sample processing. There were 148 patients tested with the Amplicor PCR and Xpert MTB/RIF assays, but only 144 had both Amplicor PCR and Xpert MTB/RIF results (31 had definite TBM [culture or microscopy positive; identification of the organism was by a niacin/nitrate test for *M. tuberculosis* and by PCR if this was negative], 60 had probable TBM, and 53 were non-TBM). Table 1 outlines the demographic and CSF characteristics that were similar and that differed in the definite TBM and non-TBM groups.

The non-TBM category ($n = 53$) comprised the following breakdown of diagnoses (number of patients): cryptococcal meningitis (29), viral meningitis (13), acute bacterial meningitis (6),

malignant meningitis (2), neurosyphilis (1), parameningeal focus (1), and other (1).

Table 2 outlines the performance outcomes of the Amplicor PCR and Xpert MTB/RIF assays. There were no differences between the Amplicor PCR and Xpert MTB/RIF performance outcomes (sensitivity, $P = 0.7$; specificity, $P = 0.6$; PPV, $P = 0.8$; and NPV, $P = 0.8$). Likelihood ratios were calculated for both the Amplicor PCR and Xpert MTB/RIF assays. The times to detection for the Amplicor PCR and Xpert MTB/RIF assays were 3.5 and 1.5 h, respectively.

There are hardly any data about the performance outcomes of newer NAATs for the diagnosis of TBM. We found no differences in the performance outcomes between the Amplicor PCR and Xpert RIF/MTB assays, and the specificities for both assays were high. The study was relatively large (31 definite TBM cases) compared to those in published reports, where the numbers of culture-confirmed CSF samples were relatively small (17–21). The sensitivity was not improved and remained suboptimal despite our use of fairly large volumes of CSF (500 μ l for the Amplicor PCR and 1,000 μ l for the Xpert MTB/RIF) compared to those in previous studies (22–26). A meta-analysis by Solomons et al. confirmed sensitivities varying from 33% to 67% (27). Previously cited reasons for the suboptimal sensitivity have included the presence of inhibitors in the CSF, paucibacillary samples below the detection limit of the assay, and the aliquot phenomenon (i.e., the initial CSF sample taken may have fewer bacilli than those taken later). We recently found that the degree of inhibition in CSF was lower than that in sputum (11). A single study comparing different aliquots of CSF using PCRs found no differences between the first,

TABLE 1 Clinical and cerebrospinal fluid data from patients with definite tuberculous meningitis (liquid culture or microscopy positive) and non-tuberculous meningitis (culture negative and no anti-TB treatment given)

| Characteristic | Results for persons with: | | P |
|--|---------------------------|-------------------------|--------|
| | Definite TBM (n = 31) | Non-TBM (n = 53) | |
| Clinical | | | |
| Mean (\pm SD) age (yr) | 32.8 (7.7) | 33.1 (11.1) | 0.9 |
| Age <36/ \geq 36 yr (no. [%]) ^a | 20/11 (64.5/35.5) | 34/19 (64.2/35.8) | 0.9 |
| Sex, male/female (no. [%]) | 15/16 (48.4/51.6) | 17/36 (32.1/67.9) | 0.2 |
| Ethnic group, BA/M/E/I (no. [%]) ^b | 31/0/0/0 (100/0/0/0) | 52/0/0/1 (98.1/0/0/1.9) | 0.9 |
| HIV status, P/N (no. [%]) ^c | 28/3 (90.3/9.7) | 47/6 (88.7/11.3) | 0.9 |
| Previous TB, yes/no/unknown (no. [%]) | 7/21/3 (22.6/67.7/9.7) | 20/31/2 (37.7/58.6/3.8) | 0.2 |
| TB contact (within 2 yr), yes/no/unknown (no. [%]) | 9/19/3 (29.0/61.3/9.7) | 14/37/2 (26.4/69.8/3.8) | 0.5 |
| Duration of illness, <6/ \geq 6 days/unknown (no. [%]) | 6/23/2 (19.4/74.2/6.5) | 7/45/1 (13.2/84.9/1.9) | 0.7 |
| Steroid treatment, yes/no (no. [%]) | 8/23 (25.8/74.2) | 10/43 (18.9/81.1) | 0.5 |
| CLAT positive, yes/no (no. [%]) | 2/27 (6.9/93.1) | 26/27 (49.1/50.9) | <0.001 |
| CD4 cells/ μ l (IQR) ^d | 116 (65–196) | 161 (78–261) | 0.1 |
| CSF parameter (median [IQR]) | | | |
| Lymphocytes (cells/ μ l) | 117 (24–242) | 32 (10–82) | 0.004 |
| Neutrophils (cells/ μ l) | 62 (24–138) | 9 (0–66) | 0.001 |
| Protein (g/liter) | 1.7 (1.2–2.5) | 1.1 (0.9–1.9) | 0.03 |
| CSF glucose (mmol/liter) | 1.1 (1.0–1.6) | 2.0 (1.5–2.7) | <0.001 |
| CSF/serum glucose ratio | 0.2 (0.2–0.3) | 0.4 (0.2–0.5) | <0.001 |
| Lymphocytes: total ratio | 0.6 (0.3–0.8) | 0.8 (0.3–1.0) | 0.2 |

^a This cut point was chosen based on criteria derived by Thwaites et al. (31).

^b BA, Black African; M, mixed race; E, European; I, Indian.

^c P, positive; N, negative.

^d IQR, interquartile range.

second, or third samples taken from the same patient (28). We recently showed that centrifugation of CSF significantly improved the Xpert MTB/RIF assay sensitivity (11). However, we were unable to ascertain whether this also applies to the Amplicor PCR assay as no concentration (centrifugation) experiments were undertaken. This study may have overestimated the sensitivity of NAATs as we used definite TBM (culture positive) as the gold standard. In the probable TBM group, the sensitivity is likely to be considerably lower because the pathogen load is often below the detection limit of the assay (11), as demonstrated in a recent large TBM study by Nhu et al. evaluating the Xpert MTB/RIF assay (29).

The specificity of the Xpert MTB/RIF assay was 94%. This is related to three patients who were classified as non-TBM (culture negative with an alternate confirmed diagnosis but Xpert MTB/RIF positive); two were categorized as having cryptococcal meningitis and one as having leukemic meningitis. Our previous work has shown that such cases (Xpert MTB/RIF positive but culture negative) are likely to be true TB positives (30), as corroborated by the high specificity obtained in large sputum-based studies where a significant minority of the patients had had previous

tuberculosis (11). It is possible that they may have had dual pathologies, but this is difficult to confirm as these patients either died or were lost to follow-up. If these culture-negative, Xpert MTB/RIF-positive persons were hypothetically designated definite TB cases, then the overall case detection rate would have improved by a further \sim 10%.

The limitations of this study include a population restricted to those who were predominantly HIV infected. This may enhance sensitivity, and CSF from such patients may theoretically harbor greater bacterial loads than CSF from immunocompetent patients. Despite the small number of culture-positive samples, this is still a relatively large cohort of definite TBM cases compared to those in similar studies.

In conclusion, this study confirms the modest but equivalent sensitivities for the Xpert MTB/RIF and Amplicor PCR assays in the absence of centrifugation in this predominantly HIV-infected cohort from a country where TB is endemic. Although the sensitivity of the Xpert MTB/RIF assay is modest, it is a useful rule-in test (diagnostic if the test is positive but does not exclude TBM if the test is negative) and thus the Xpert MTB/RIF is useful for the

TABLE 2 Diagnostic accuracy of Amplicor PCR and Xpert MTB/RIF tests using liquid culture and smear microscopy as the reference standard

| Test | Performance outcome (95% CI) | | | | | LR ^a | |
|---------------|---|--|---------------------------------|---------------------------------|----------------------------------|-----------------|-----------------|
| | Sensitivity | Specificity | PPV | NPV | Agreement | LR ⁺ | LR ⁻ |
| Amplicor | 46 (31, 60) ^b 21/46 ^d | 99 (93, 100) ^c 80/81 ^d | 96 (77, 100) 21/22 ^d | 76 (67, 84) 80/105 ^d | 80 (72, 86) 101/127 ^d | 37.0 | -0.55 |
| Xpert MTB/RIF | 50 ¹ (33, 67) 18/36 ^d | 94 ² (84, 99) 50/53 ^d | 86 (64, 97) 18/21 ^d | 74 (61, 84) 50/68 ^d | 76 (66, 85) 68/89 ^d | 8.8 | -0.53 |

^a LR, likelihood ratio. LR⁺ is the ratio of the probability of a positive test among the truly positive subjects to the probability of a positive test among the truly negative, and LR⁻ is the ratio of the probability of a negative test among the truly positive subjects to the probability of a negative test among the truly negative subjects.

^b Comparison of sensitivity between Amplicor and Xpert MTB/RIF tests: $P = 0.7$.

^c comparison of specificity between Amplicor and Xpert MTB/RIF tests: $P = 0.6$.

^d An explanation of this fraction is included in the supplementary material.

rapid diagnosis of TBM where delay may otherwise result in excess mortality and significant morbidity. However, studies in other settings, in HIV-uninfected populations, and with variations in processing methods (centrifugation, CSF volume, etc) are now required to improve the sensitivity.

ACKNOWLEDGMENTS

We are extremely grateful to the patients, registrars in the department of neurology, and nurses for facilitating this study. We are grateful to the Province of KwaZulu-Natal and the TB program for facilitating the study.

V.B.P. and K.D. originated the study and wrote the paper; K.D. and T.N. supervised the study; R.S. did the laboratory work; L.L., B.M., and G.T. assisted with the laboratory work; and C.C. was the statistician responsible for the statistical analysis.

This work was supported by the Columbia University-Southern African Fogarty AIDS International Training and Research Program funded by the Fogarty International Centre, the National Institutes of Health (grant D43TW00231 to V.B.P.), a South African MRC grant (V.B.P. and K.D.), the EU FP7 programme (TB susgent) (V.B.P. and K.D.), the South African NRF Research Chairs Initiative (SARChI) (T.N. and K.D.), an SA MRC Career Development Award (K.D.), and the EDCTP (TESA and TB-NEAT). Publication of this article has been made possible through support from the Victor Daitz Information Gateway, an initiative of the Victor Daitz Foundation and the University of KwaZulu-Natal. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

We declare no conflicts of interest.

REFERENCES

- World Health Organization. 2011. Global tuberculosis control 2011. World Health Organization, Geneva, Switzerland.
- Braunwald E, Fauci AS, Kasper DL, Hauser SL, Longo DL, Jameson JL. 2001. Harrison's principles of internal medicine, 15th ed, vol 1. McGraw-Hill, New York, NY.
- Ozbay B, Uzun K. 2002. Extrapulmonary tuberculosis in high prevalence of tuberculosis and low prevalence of HIV. *Clin. Chest Med.* 23:351–354. [http://dx.doi.org/10.1016/S0272-5231\(02\)00002-3](http://dx.doi.org/10.1016/S0272-5231(02)00002-3).
- Garcia-Monco JC. 1999. Central nervous system tuberculosis. *Neurol. Clin.* 17:737–759. [http://dx.doi.org/10.1016/S0733-8619\(05\)70164-X](http://dx.doi.org/10.1016/S0733-8619(05)70164-X).
- Karsteadt AS, Valtachanova S, Barriere R, Crewe-Brown HH. 1998. Tuberculous meningitis in South African urban adults. *Q. J. M.* 91:743–747. <http://dx.doi.org/10.1093/qjmed/91.11.743>.
- Rock RB, Olin M, Baker CA, Molitor TW, Peterson PK. 2008. Central nervous system tuberculosis: pathogenesis and clinical aspects. *Clin. Microbiol. Rev.* 21:243–261. <http://dx.doi.org/10.1128/CMR.00042-07>.
- Vinnard C, Macgregor RR. 2009. Tuberculous meningitis in HIV-infected individuals. *Curr. HIV/AIDS Rep.* 6:139–145. <http://dx.doi.org/10.1007/s11904-009-0019-7>.
- Pai M, Flores LL, Pai N, Hubbard A, Riley LW, Colford JM, Jr. 2003. Diagnostic accuracy of nucleic acid amplification tests for tuberculous meningitis: a systematic review and meta-analysis. *Lancet Infect. Dis.* 3:633–643. [http://dx.doi.org/10.1016/S1473-3099\(03\)00772-2](http://dx.doi.org/10.1016/S1473-3099(03)00772-2).
- Lawn SD, Nicol MP. 2011. Xpert MTB/RIF assay: development, evaluation and implementation of a new rapid molecular diagnostic for tuberculosis and rifampicin resistance. *Future Microbiol.* 6:1067–1082. <http://dx.doi.org/10.2217/fmb.11.84>.
- Reischl U, Lehn N, Wolf H, Naumann L. 1998. Clinical evaluation of the automated COBAS Amplicor MTB assay for testing respiratory and non-respiratory specimens. *J. Clin. Microbiol.* 36:2853–2860.
- Patel VB, Theron G, Lenders L, Matinyena B, Connolly C, Singh R, Coovadia Y, Ndung'u T, Dheda K. 2013. Diagnostic accuracy of quantitative PCR (Xpert MTB/RIF) for tuberculous meningitis in a high burden setting: a prospective study. *PLoS Med.* 10:e1001536. <http://dx.doi.org/10.1371/journal.pmed.1001536>.
- Patel VB, Singh R, Connolly C, Coovadia Y, Peer AK, Parag P, Kasprovicz V, Zumla A, Ndung'u T, Dheda K. 2010. Cerebrospinal T-cell responses aid in the diagnosis of tuberculous meningitis in a human immunodeficiency virus- and tuberculosis-endemic population. *Am. J. Respir. Crit. Care Med.* 182:569–577. <http://dx.doi.org/10.1164/rccm.200912-1931OC>.
- Cepheid. 2009. Xpert MTB/RIF package insert 300-7810 revision A. Cepheid, Sunnyvale, CA.
- Marais S, Thwaites G, Schoeman JF, Torok ME, Misra UK, Prasad K, Donald PR, Wilkinson RJ, Marais BJ. 2010. Tuberculous meningitis: a uniform case definition for use in clinical research. *Lancet Infect. Dis.* 10:803–812. [http://dx.doi.org/10.1016/S1473-3099\(10\)70138-9](http://dx.doi.org/10.1016/S1473-3099(10)70138-9).
- Thwaites G, Fisher M, Hemingway C, Scott G, Solomon T, Innes J. 2009. British Infection Society guidelines for the diagnosis and treatment of tuberculosis of the central nervous system in adults and children. *J. Infect.* 59:167–187. <http://dx.doi.org/10.1016/j.jinf.2009.06.011>.
- Thwaites G, Chau TT, Mai NT, Drobniowski F, McAdam K, Farrar J. 2000. Tuberculous meningitis. *J. Neurol. Neurosurg. Psychiatry* 68:289–299. <http://dx.doi.org/10.1136/jnnp.68.3.289>.
- Causse M, Ruiz P, Gutierrez-Aroca JB, Casal M. 2011. Comparison of two molecular methods for rapid diagnosis of extrapulmonary tuberculosis. *J. Clin. Microbiol.* 49:3065–3067. <http://dx.doi.org/10.1128/JCM.00491-11>.
- Ehlers S, Ignatius R, Regnath T, Hahn H. 1996. Diagnosis of extrapulmonary tuberculosis by Gen-Probe amplified *Mycobacterium tuberculosis* direct test. *J. Clin. Microbiol.* 34:2275–2279.
- Eing BR, Becker A, Sohns A, Ringelmann R. 1998. Comparison of Roche Cobas Amplicor *Mycobacterium tuberculosis* assay with in-house PCR and culture for detection of *M. tuberculosis*. *J. Clin. Microbiol.* 36:2023–2029.
- Jönsson B, Ridell M. 2003. The Cobas Amplicor MTB test for detection of *Mycobacterium tuberculosis* complex from respiratory and non-respiratory clinical specimens. *Scand. J. Infect. Dis.* 35:372–377. <http://dx.doi.org/10.1080/00365540310012244>.
- Tortoli E, Russo C, Piersimoni C, Mazzola E, Dal MP, Pascarella M, Borroni E, Mondo A, Piana F, Scarparo C, Coltella L, Lombardi G, Cirillo DM. 2012. Clinical validation of Xpert MTB/RIF for the diagnosis of extrapulmonary tuberculosis. *Eur. Respir. J.* 40:442–447. <http://dx.doi.org/10.1183/09031936.00176311>.
- Armand S, Vanhuls P, Delcroix G, Courcol R, Lemaitre N. 2011. Comparison of the Xpert MTB/RIF test with an IS6110-TaqMan real-time PCR assay for direct detection of *Mycobacterium tuberculosis* in respiratory and nonrespiratory specimens. *J. Clin. Microbiol.* 49:1772–1776. <http://dx.doi.org/10.1128/JCM.02157-10>.
- Bonington A, Strang JI, Klapper PE, Hood SV, Rubombora W, Penny M, Willers R, Wilkins EG. 1998. Use of Roche Amplicor *Mycobacterium tuberculosis* PCR in early diagnosis of tuberculous meningitis. *J. Clin. Microbiol.* 36:1251–1254.
- Caws M, Wilson SM, Clough C, Drobniowski F. 2000. Role of IS6110-targeted PCR, culture, biochemical, clinical, and immunological criteria for diagnosis of tuberculous meningitis. *J. Clin. Microbiol.* 38:3150–3155. <http://jcm.asm.org/content/38/9/3150.long>.
- Daley P, Thomas S, Pai M. 2007. Nucleic acid amplification tests for the diagnosis of tuberculous lymphadenitis: a systematic review. *Int. J. Tuberc. Lung Dis.* 11:1166–1176. <http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0024738/>.
- Deshpande PS, Kashyap RS, Ramteke SS, Nagdev KJ, Purohit HJ, Taori GM, Dagainawala HF. 2007. Evaluation of the IS6110 PCR assay for the rapid diagnosis of tuberculous meningitis. *Cerebrospinal Fluid Res.* 4:10. <http://dx.doi.org/10.1186/1743-8454-4-10>.
- Solomons RS, van Elsland SL, Visser DH, Hoek KG, Marais BJ, Schoeman JF, van Furth AM. 2014. Commercial nucleic acid amplification tests in tuberculous meningitis—a meta-analysis. *Diagn. Microbiol. Infect. Dis.* 78:398–403. <http://dx.doi.org/10.1016/j.diagmicrobio.2014.01.002>.
- Bhigjee AI, Padayachee R, Paruk H, Hallwirth-Pillay KD, Marais S, Connolly C. 2007. Diagnosis of tuberculous meningitis: clinical and laboratory parameters. *Int. J. Infect. Dis.* 11:348–354. <http://dx.doi.org/10.1016/j.ijid.2006.07.007>.
- Nhu NT, Heemskerck D, Thu do DA, Chau TT, Mai NT, Nghia HD, Loc PP, Ha DT, Merson L, Thinh TT, Day J, Chau Nv, Wolbers M, Farrar J, Caws M. 2014. Evaluation of GeneXpert MTB/RIF for diagnosis of tuberculous meningitis. *J. Clin. Microbiol.* 52:226–233. <http://dx.doi.org/10.1128/JCM.01834-13>.
- Theron G, Peter J, van Zyl-Smit R, Mishra H, Streicher E, Murray S, Dawson R, Whitelaw A, Hoelscher M, Sharma S, Pai M, Warren R, Dheda K. 2011. Evaluation of the Xpert MTB/RIF assay for the diagnosis of pulmonary tuberculosis in a high HIV prevalence setting. *Am. J. Respir. Crit. Care Med.* 184:132–140. <http://dx.doi.org/10.1164/rccm.201101-0056OC>.
- Thwaites GE, Chau TT, Stepniowska K, Phu NH, Chuong LV, Sinh DX, White NJ, Parry CM, Farrar JJ. 2002. Diagnosis of adult tuberculous meningitis by use of clinical and laboratory features. *Lancet* 360:1287–1292. [http://dx.doi.org/10.1016/S0140-6736\(02\)11318-3](http://dx.doi.org/10.1016/S0140-6736(02)11318-3).