




Brief report: International perspectives on the pediatric COVID#19 experience

Item Type	Article
Authors	Yilmaz, O.;Gochicoa-Rangel, L.;Blau, H.;Epaud, R.;Lands, L.C.;Lombardi, E.;Moore, P.E.;Stein, R.T.;Wong, G.W.K.;Zar, H.J.
Citation	Yilmaz O, Gochicoa-Rangel L, Blau H, Epaud R, Lands LC, Lombardi E, Moore PE, Stein RT, Wong GWK, Zar HJ. Brief report: International perspectives on the pediatric COVID-19 experience. <i>Pediatr Pulmonol.</i> 2020 Jul;55(7):1598-1600. doi: 10.1002/ppul.24800.
DOI	10.1002/ppul.24800
Publisher	Wiley
Journal	Pediatric Pulmonology
Rights	Attribution 3.0 United States
Download date	2025-04-28 18:58:00
Item License	http://creativecommons.org/licenses/by/3.0/us/
Link to Item	https://pubmed.ncbi.nlm.nih.gov/32356583/

COMMENTARY

Brief report: International perspectives on the pediatric COVID-19 experience

Ozge Yilmaz MD¹  | Laura Gochicoa-Rangel MD² | Hannah Blau MBBS³ |
Ralph Epaud MD⁴  | Larry C. Lands MD, PhD⁵ | Enrico Lombardi MD⁶ |
Paul E. Moore MD⁷  | Renato T. Stein MD⁸ | Gary W. K. Wong MD⁹ |
Heather J. Zar MD, PhD¹⁰

¹Department of Pediatric Allergy and Pulmonology, Medical Faculty, Celal Bayar University, Manisa, Turkey

²Department of Pulmonary Physiology, National Institute of Respiratory Diseases, Mexico City, Mexico

³Pulmonary Institute at Schneider Children's Medical Center, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

⁴Department of Pediatrics, CHI Créteil, Service de Pédiatrie, INSERM U955 (IMRB), équipe GEIC2O, UPEC, Créteil, France

⁵Department of Pediatrics, Montreal Children's Hospital, McGill University Health Centre, Montreal, Quebec, Canada

⁶Pediatric Pulmonary Unit, Meyer Children's Hospital, Florence, Italy

⁷Department of Pediatrics, Monroe Carell Jr. Children's Hospital, Vanderbilt University Medical Center, Nashville, Tennessee

⁸Pontifical Catholic University of Rio Grande do Sul, Porto Alegre, Brazil

⁹Department of Pediatrics, Prince of Wales Hospital, Chinese University of Hong Kong, Hong Kong

¹⁰Department of Paediatrics and Child Health, SA MRC Unit on Child and Adolescent Health, Red Cross Children's Hospital, University of Cape Town, Cape Town, South Africa

Correspondence

Ozge Yilmaz, MD, Department of Pediatric Allergy and Pulmonology, Celal Bayar University, 45030, Manisa, Turkey.
Email: oyilmaz_76@hotmail.com

Laura Gochicoa-Rangel, MD, National Institute of Respiratory Diseases, Mexico City, Mexico.
Email: drgochis@gmail.com

Abstract

The 2019 novel coronavirus (SARS-CoV-2) is endangering human health worldwide; scarcity of published pediatric cases and current literature and the absence of evidence-based guidelines necessitate international sharing of experience and personal communication. On 31 March 2020 the International Committee of the American Thoracic Society Pediatrics Assembly recorded an online podcast, during which pediatric pulmonologists worldwide shared their experience on the novel coronavirus disease (COVID-19) in children. The aim was to share personal experience in organizing pediatric care in different health care settings globally, protecting health care workers, and isolation practices. This manuscript summarizes the common themes of the podcast which centered around three main topics: more benign clinical disease and progression in pediatric cases compared to adults, a strong need for strategies to protect health care workers, and social or economic disparities as a barrier to successful pandemic control.

KEYWORDS

COVID-19, disparity, pandemic, pediatric pulmonology

On 31 March 2020 the International Committee of the American Thoracic Society Pediatrics Assembly recorded an online podcast, during which pediatric pulmonologists worldwide shared their

experience on the novel coronavirus disease (COVID-19) in children. The aim was not to share centralized information about pediatric patient numbers or treatment protocols which is available through

much international epidemiology and health care websites; but to share personal experience in organizing pediatric care in different health care settings globally, protecting health care workers, and isolation practices. The 2019 novel coronavirus (SARS-CoV-2) is endangering human health worldwide; scarcity of published pediatric cases and current literature and the absence of evidence-based guidelines necessitate international sharing of experience and personal communication.^{1,2}

The common themes of the podcast centered around three main topics: more benign clinical disease and progression in pediatric cases compared with adults, a strong need for strategies to protect health care workers, and social or economic disparities as a barrier to successful pandemic control.

1 | BENIGN COURSE OF PEDIATRIC CASES

As depicted in recent articles from China and Korea, pediatric cases constitute a small percentage of the total COVID-19 hospitalized cases worldwide with low mortality being reported in children.¹⁻³ Evaluation of the symptomatic and asymptomatic children younger than 16 years of age in Wuhan with known contact with COVID-19 cases found that 12.3% of those tested were confirmed to have SARS-CoV-2 infection; pneumonia was the most common diagnosis amongst symptomatic children.⁴ Similarly, the Center for Disease Control and prevention in China reported that only 2% of hospitalized cases were younger than 19 years of age; in Korea under 5% of cases were under 19 years of age.^{3,5} Fever was not a prominent finding, could occur in around 50% while radiologically ground-glass opacities were commonly encountered. The majority of the children had milder disease and were discharged from the hospital.¹ Moreover, SARS-CoV-2 infection has a wide range of symptoms in children including diarrhea, vomiting, and fatigue.⁴ Therefore, children play an important role in spreading of the infection in the community due to the asymptomatic nature of the infection, making extensive testing and isolation of children vital for infection control.

The local experience of the podcast participants was in line with these publications in terms of frequency and severity. Children with severe clinical presentation usually had underlying comorbid diseases such as oncologic malignancies, diabetes, and immunodeficiency. The chronic respiratory disease did not appear to be emerging as a risk factor for severe COVID-19 disease; however, long term follow-up is needed to confirm this observation. Lack of information on long term prognosis of asymptomatic children and those with pneumonia should be a research priority.

2 | SOCIAL DISTANCING MEASURES AND CLINICAL SERVICES

Pediatric care in many hospitals is organized to separate patients with suspected or confirmed COVID-19 from other patients. Emergency departments have separated tracks to avoid contact of

suspected COVID-19 patients, who are admitted to a different part of the building. Similarly, wards for hospitalization are separated for suspected and confirmed COVID-19 cases, with separate staff in emergency and inpatients areas.

In many hospitals, elective ambulatory pediatric care, as well as lung function testing, is halted in favor of social distancing measures. Moreover, telemedicine measures are implemented or starting to be implemented in many parts of the world, especially for children with chronic diseases. Telemedicine is used both for consultation by the patients and to keep contact with families and children with chronic diseases. Another concern is the care of children with other diseases. Globally, ambulatory clinics and elective procedures have been curtailed. The availability of urgent health care to children with non-COVID diseases is vital to protect child health. As the number of pediatric COVID cases is low and the ambulatory activities have been drastically reduced, many institutions have pediatricians or health care personnel normally working in the pediatric department to help adult colleagues.

One issue for pediatric care in COVID-19 patients is nebulization practice. Nebulized treatment should be avoided and replaced with inhaled treatment with spacers and meter-dosed inhalers. Nebulizers may carry the potential for the spread of infection due to aerosols generated, whereas spacers are much easier for sterilization issues. Similarly, high flow oxygen should be avoided.

Social distancing seems to be the major tool to prevent the spread of the infection. Although asymptomatic, children may shed the virus especially amongst household members thus separating the elderly from young children is important. However, social disparities will be the major obstacle to implement social distancing measures in many countries.

3 | SAFETY OF HEALTH CARE WORKERS

The safety of health care workers is a major concern raised by all participants. Asymptomatic or mildly symptomatic infections represent an important risk of spreading the infection to health care workers. Some hospitals take measures such as daily fever checks for all those who enter the hospital, including health care workers, and COVID-19 testing for all children requiring hospitalization, even if they do not have respiratory symptoms. Closed-circuit television and online care of patients with COVID-19 was implemented in some institutions to decrease contact of health care workers with the infected subjects. Moreover, health care workers taking care of potentially infected children should perform proper hand hygiene and use of appropriate personal protective equipment (PPE), including use of a respirator, gowns, gloves, caps, and eye protection.^{6,7}

There are different practices in terms of the use of masks for health care workers. In some institutions, all hospital workers, including physicians, nurses, and administrative personnel, are using masks, and in some only those working with patients are using masks. The use of universal masks by the public and in health care facilities may reduce transmission of COVID-19 and other respiratory viruses, thus decreasing the burden of respiratory infections on health care facilities.⁸

Availability of surgical and N95 masks is an issue, and these should be reserved for use in health care facilities given the global shortages. Homemade cloth masks are appropriate for use by the public.

Anxiety, stress, and exhaustion among health care workers are also key concerns.

A flexible action plan is important component management. Action plan on quarantine measures, separating health care workers who have contact, and who do not have contact with patients is important. The flexibility of the action plan to move adequate health care workers to the required areas in the hospital will be required during the pandemic.

As the epidemic moves from high-income countries to Africa and Latin America, key concerns about access to PPE for HCWs, the ability of health systems to address the burden, the inability to undertake social distancing and frequent hand washing in poor communities, and potential impact on the severity of disease by highly prevalent diseases like TB or HIV are pressing. Further the impact of the epidemic and on lockdowns on economies and on poverty remain critical issues for child health.

ACKNOWLEDGMENT

This report was based on the podcast recorded by the American Thoracic Society. All authors contributed equally so all needs to get credit as first authors.

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

AUTHOR CONTRIBUTION

All authors have contributed equally to the content and preparation of the manuscript, therefore, would like to receive credit as first authors.

ORCID

Ozge Yilmaz  <http://orcid.org/0000-0001-6051-5020>

Ralph Epaud  <http://orcid.org/0000-0003-3830-1039>

Paul E. Moore  <http://orcid.org/0000-0001-5123-207X>

REFERENCES

1. Wang Y, Wang Y, Chen Y, Qin Q. Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID-19) implicate special control measures. *J Med Virol.* 2020;92:568-576.
2. Bedford J, Enria D, Giesecke J, et al. COVID-19: towards controlling of a pandemic. *Lancet.* 2020;395(10229):1015-1018.
3. Korean Society of Infectious Diseases, Korean Society of Pediatric Infectious Diseases, Korean Society of Epidemiology, Korean Society for Antimicrobial Therapy, Korean Society for Healthcare-associated Infection Control and Prevention, Korea Centers for Disease Control and Prevention. Report on the Epidemiological Features of Coronavirus Disease 2019 (COVID-19) Outbreak in the Republic of Korea from January 19 to March 2, 2020. *J Korean Med Sci.* 2020;35:e112.
4. Lu X, Zhang L, Du H, et al. SARS-CoV-2 infection in children. *N Engl J Med.* 2020;382:1663-1665.
5. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA.* 2020;323:1239-1242.
6. World Health Organization. Infection prevention and control during health care when novel coronavirus (nCoV) infection is suspected; 2020. [https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-\(ncov\)-infection-is-suspected-20200125](https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-(ncov)-infection-is-suspected-20200125). Accessed April 15, 2020.
7. Centers for Disease Control and Prevention. Interim infection prevention and control recommendations for patients with suspected or confirmed coronavirus disease 2019 (COVID-19) in healthcare settings; 2020. <https://www.cdc.gov/coronavirus/2019-ncov/infection-control/control-recommendations.html>. Accessed April 15, 2020.
8. Klompas M, Morris CA, Sinclair J, Pearson M, Shenoy ES. Universal masking in hospitals in the Covid-19 era. *New Engl J Med.* 2020.

How to cite this article: Yilmaz O, Gochicoa-Rangel L, Blau H, et al. Brief report: International perspectives on the pediatric COVID-19 experience. *Pediatric Pulmonology.* 2020;55:1598-1600. <https://doi.org/10.1002/ppul.24800>